PROOF OF A BASIC HYPERGEOMETRIC SUPERCONGRUENCE MODULO THE FIFTH POWER OF A CYCLOTOMIC POLYNOMIAL

VICTOR J. W. GUO AND MICHAEL J. SCHLOSSER

Abstract. By means of the q-Zeilberger algorithm, we prove a basic hypergeometric supercongruence modulo the fifth power of the cyclotomic polynomial $\Phi_n(q)$. This result appears to be quite unique, as in the existing literature so far no basic hypergeometric supercongruences modulo a power greater than the fourth of a cyclotomic polynomial have been proved. We also establish a couple of related results, including a parametric supercongruence.

1. Introduction

In 1997, Van Hamme [27] conjectured that 13 Ramanujan-type series including

$$\sum_{k=0}^{\infty}(-1)^k(4k+1)\frac{(\frac{1}{2})_k^3}{k!^3}=\frac{2}{\pi},$$

admit nice p-adic analogues, such as

$$\sum_{k=0}^{\frac{p-1}{2}}(-1)^k(4k+1)\frac{(\frac{1}{2})_k^3}{k!^3}\equiv p\,(-1)^{\frac{p-1}{2}} \pmod{p^3},$$

where $(a)_n=a(a+1)\cdots(a+n-1)$ denotes the Pochhammer symbol and p is an odd prime. Up to present, all of the 13 supercongruences have been confirmed. See [21,24] for historic remarks on these supercongruences. Recently, q-analogues of congruences and supercongruences have caught the interests of many authors (see, for example, [1–20,25,26,29]). In particular, the first author and Zudilin [16] devised a method, called 'creative microscoping', to prove quite a few q-supercongruences by introducing an additional parameter a. In [13], the authors of the present paper proved many additional q-supercongruences by the creative microscoping method. Supercongruences modulo a higher integer power of a prime, or, in the q-case, of a cyclotomic polynomial, are very special and usually difficult to prove. As far as we know, until now the result

$$\sum_{k=0}^{\frac{n-1}{2}}[4k+1]\frac{(q;q^2)_k^4}{(q^2;q^2)_k^4}\equiv q^{\frac{1-n}{2}}[n]+\frac{(n^2-1)(1-q)^2}{24}q^{\frac{1-n}{2}}[n]^3 \pmod{[n]\Phi_n(q)^3}, \tag{1}$$

2010 *Mathematics Subject Classification.* Primary 33D15; Secondary 11A07, 11F33.

Key words and phrases. basic hypergeometric series, q-series, supercongruences, identities.

The first author was partially supported by the National Natural Science Foundation of China (grant 11771175).

for an odd positive integer n, due to the first author and Wang [15], is the unique q-supercongruence modulo $[n]\Phi_n(q)^3$ in the literature that was completely proved. (Several similar conjectural q-supercongruences are stated in [13] and in [16].) The purpose of this paper is to establish an even higher q-congruence, namely modulo a fifth power of a cyclotomic polynomial. Specifically, we prove the following three theorems. (The first two together confirm a conjecture by the authors [13, Conjecture 5.4]).

Theorem 1.1. *Let $n>1$ be a positive odd integer. Then*

$$\sum_{k=0}^{\frac{n+1}{2}}[4k-1]\frac{(q^{-1};q^2)_k^4}{(q^2;q^2)_k^4}q^{4k}\equiv -(1+3q+q^2)[n]^4 \pmod{[n]^4\Phi_n(q)}, \tag{2a}$$

and

$$\sum_{k=0}^{n-1}[4k-1]\frac{(q^{-1};q^2)_k^4}{(q^2;q^2)_k^4}q^{4k}\equiv -(1+3q+q^2)[n]^4 \pmod{[n]^4\Phi_n(q)}. \tag{2b}$$

Theorem 1.2. *Let $n>1$ be a positive odd integer. Then*

$$\sum_{k=0}^{\frac{n+1}{2}}[4k-1]\frac{(aq^{-1};q^2)_k(q^{-1}/a;q^2)_k(q^{-1};q^2)_k^2}{(aq^2;q^2)_k(q^2/a;q^2)_k(q^2;q^2)_k^2}q^{4k}\equiv 0 \pmod{[n]^2(1-aq^n)(a-q^n)},$$

and

$$\sum_{k=0}^{n-1}[4k-1]\frac{(aq^{-1};q^2)_k(q^{-1}/a;q^2)_k(q^{-1};q^2)_k^2}{(aq^2;q^2)_k(q^2/a;q^2)_k(q^2;q^2)_k^2}q^{4k}\equiv 0 \pmod{[n]^2(1-aq^n)(a-q^n)}.$$

The $a=-1$ case of Theorem 1.2 admits an even stronger q-congruence.

Theorem 1.3. *Let $n>1$ be a positive odd integer. Then*

$$\sum_{k=0}^{\frac{n+1}{2}}[4k-1]\frac{(q^{-2};q^4)_k^2}{(q^4;q^4)_k^2}q^{4k}\equiv -q^n(1-q+q^2)[n]_{q^2}^2 \pmod{[n]_{q^2}^2\Phi_n(q^2)}, \tag{3a}$$

and

$$\sum_{k=0}^{n-1}[4k-1]\frac{(q^{-2};q^4)_k^2}{(q^4;q^4)_k^2}q^{4k}\equiv -(1-q+q^2)[n]_{q^2}^2 \pmod{[n]_{q^2}^2\Phi_n(q^2)}. \tag{3b}$$

In the above q-supercongruences and in what follows,

$$(a;q)_n=(1-a)(1-aq)\cdots(1-aq^{n-1})$$

is the q-shifted factorial,

$$[n]=[n]_q=1+q+\cdots+q^{n-1}$$

is the q-number,

$$\begin{bmatrix}n\\k\end{bmatrix}=\begin{bmatrix}n\\k\end{bmatrix}_q:=\frac{(q;q)_n}{(q;q)_k(q;q)_{n-k}}$$

is the q-binomial coefficient, and $\Phi_n(q)$ is the n-th cyclotomic polynomial of q. Note that the congruences in Theorem 1.1 modulo $[n]\Phi_n(q)^2$ and the congruences in Theorem 1.2 modulo $[n](1-aq^n)(a-q^n)$ have already been proved by the authors in [13, eqs. (5.5) and (5.10)].

2. Proof of Theorem 1.1 by the Zeilberger algorithm

The Zeilberger algorithm (cf. [22]) can be used to find that the functions

$$f(n,k)=(-1)^k\frac{(4n-1)(-\frac{1}{2})_n^3(-\frac{1}{2})_{n+k}}{(1)_n^3(1)_{n-k}(-\frac{1}{2})_k^2},$$

$$g(n,k)=(-1)^{k-1}\frac{4(-\frac{1}{2})_n^3(-\frac{1}{2})_{n+k-1}}{(1)_{n-1}^3(1)_{n-k}(-\frac{1}{2})_k^2}$$

satisfy the relation

$$(2k-3)f(n,k-1)-(2k-4)f(n,k)=g(n+1,k)-g(n,k).$$

Of course, given this relation, it is not difficult to verify by hand that it is satisfied by the above pair of doubly-indexed sequences $f(n,k)$ and $g(n,k)$.

Here we use the convention $1/(1)_m=0$ for all negative integers m. We now define the q-analogues of $f(n,k)$ and $g(n,k)$ as follows:

$$F(n,k)=(-1)^kq^{(k-2)(k-2n+1)}\frac{[4n-1](q^{-1};q^2)_n^3(q^{-1};q^2)_{n+k}}{(q^2;q^2)_n^3(q^2;q^2)_{n-k}(q^{-1};q^2)_k^2},$$

$$G(n,k)=\frac{(-1)^{k-1}q^{(k-2)(k-2n+3)}(q^{-1};q^2)_n^3(q^{-1};q^2)_{n+k-1}}{(1-q)^2(q^2;q^2)_{n-1}^3(q^2;q^2)_{n-k}(q^{-1};q^2)_k^2},$$

where we have used the convention that $1/(q^2;q^2)_m=0$ for $m=-1,-2,\ldots$. Then the functions $F(n,k)$ and $G(n,k)$ satisfy the relation

$$[2k-3]F(n,k-1)-[2k-4]F(n,k)=G(n+1,k)-G(n,k). \tag{4}$$

Indeed, it is straightforward to obtain the following expressions:

$$\frac{F(n,k-1)}{G(n,k)}=\frac{q^{2n-4k+6}(1-q)(1-q^{4n-1})(1-q^{2k-3})^2}{(1-q^{2n-2k+2})(1-q^{2n})^3},$$

$$\frac{F(n,k)}{G(n,k)}=-\frac{q^{4-2k}(1-q)(1-q^{4n-1})(1-q^{2n+2k-3})}{(1-q^{2n})^3},$$

$$\frac{G(n+1,k)}{G(n,k)}=\frac{q^{4-2k}(1-q^{2n-1})^3(1-q^{2n+2k-3})}{(1-q^{2n})^3(1-q^{2n-2k+2})}.$$

It is easy to verify the identity

$$\frac{q^{2n-4k+6}(1-q^{4n-1})(1-q^{2k-3})^3}{(1-q^{2n-2k+2})(1-q^{2n})^3}+\frac{q^{4-2k}(1-q^{2k-4})(1-q^{4n-1})(1-q^{2n+2k-3})}{(1-q^{2n})^3}$$
$$=\frac{q^{4-2k}(1-q^{2n-1})^3(1-q^{2n+2k-3})}{(1-q^{2n})^3(1-q^{2n-2k+2})}-1,$$

which is equivalent to (4). (Alternatively, we could have established (4) by only guessing $F(n,k)$ and invoking the q-Zeilberger algorithm [28].)

Let $m>1$ be an odd integer. Summing (4) over n from 0 to $(m+1)/2$, we get

$$[2k-3]\sum_{n=0}^{\frac{m+1}{2}}F(n,k-1)-[2k-4]\sum_{n=0}^{\frac{m+1}{2}}F(n,k)=G\left(\frac{m+3}{2},k\right)-G(0,k)$$
$$=G\left(\frac{m+3}{2},k\right). \tag{5}$$

We readily compute

$$G\left(\frac{m+3}{2},1\right)=\frac{q^{m-1}(q^{-1};q^2)^4_{(m+3)/2}}{(1-q)^2(q^2;q^2)^4_{(m+1)/2}(1-q^{-1})^2}$$
$$=\frac{q^{m-3}[m]^4}{[m+1]^4(-q;q)^8_{(m-1)/2}}\begin{bmatrix}m-1\\(m-1)/2\end{bmatrix}^4, \tag{6a}$$

and

$$G\left(\frac{m+3}{2},2\right)=-\frac{(q^{-1};q^2)^3_{(m+3)/2}(q^{-1};q^2)_{(m+5)/2}}{(1-q)^2(q^2;q^2)^3_{(m+1)/2}(q^2;q^2)_{(m-1)/2}(q^{-1};q^2)^2_2}$$
$$=-\frac{q^{-2}[m]^4[m+2]}{[m+1]^3(-q;q)^8_{(m-1)/2}}\begin{bmatrix}m-1\\(m-1)/2\end{bmatrix}^4. \tag{6b}$$

Combining (5) and (6), we have

$$\sum_{n=0}^{\frac{m+1}{2}}F(n,0)=\frac{[-2]}{[-1]}\sum_{n=0}^{\frac{m+1}{2}}F(n,1)+\frac{1}{[-1]}G\left(\frac{m+3}{2},1\right)$$
$$=\frac{1+q}{q}G\left(\frac{m+3}{2},2\right)-qG\left(\frac{m+3}{2},1\right)$$
$$=-\frac{(1+q)[m]^4[m+1][m+2]+q^{m+1}[m]^4}{q^3[m+1]^4(-q;q)^8_{(m-1)/2}}\begin{bmatrix}m-1\\(m-1)/2\end{bmatrix}^4,$$

i.e.,

$$\sum_{n=0}^{\frac{m+1}{2}}[4n-1]\frac{(q^{-1};q^2)_n^4}{(q^2;q^2)_n^4}q^{4n}=-\frac{(1+q)[m]^4[m+1][m+2]+q^{m+1}[m]^4}{q[m+1]^4(-q;q)_{(m-1)/2}^8}\begin{bmatrix}m-1\\(m-1)/2\end{bmatrix}^4. \quad (7)$$

By [4, Lemma 2.1] (or [3, Lemma 2.1]), we have $(-q;q)_{(m-1)/2}^2\equiv q^{(m^2-1)/8} \pmod{\Phi_m(q)}$. Moreover, it is easy to see that

$$\begin{bmatrix}m-1\\(m-1)/2\end{bmatrix}=\prod_{k=1}^{(m-1)/2}\frac{1-q^{m-k}}{1-q^k}\equiv\prod_{k=1}^{(m-1)/2}\frac{1-q^{-k}}{1-q^k}=(-1)^{(m-1)/2}q^{(1-m^2)/8} \pmod{\Phi_m(q)},$$

and $[m]$ is relatively prime to $(-q;q)_{(m-1)/2}$. It follows from (7) that

$$\sum_{n=0}^{\frac{m+1}{2}}[4n-1]\frac{(q^{-1};q^2)_n^4}{(q^2;q^2)_n^4}q^{4n}\equiv-((1+q)^2+q)[m]^4 \pmod{[m]^4\Phi_m(q)}.$$

Concluding, the congruence (2a) holds.

Similarly, summing (4) over n from 0 to $m-1$, we get

$$[2k-3]\sum_{n=0}^{m-1}F(n,k-1)-[2k-4]\sum_{n=0}^{m-1}F(n,k)=G(m,k),$$

and so

$$\begin{aligned}\sum_{n=0}^{m-1}[4n-1]\frac{(q^{-1};q^2)_n^4}{(q^2;q^2)_n^4}q^{4n}&=\frac{1+q}{q}G(m,2)-qG(m,1)\\&=-\frac{(1+q)[2m-2][2m-1]+q^{2m-2}}{q(-q;q)_{m-1}^8}\begin{bmatrix}2m-2\\m-1\end{bmatrix}^4. \quad (8)\end{aligned}$$

It is easy to see that

$$\frac{1}{[m]}\begin{bmatrix}2m-2\\m-1\end{bmatrix}=\frac{1}{[m-1]}\begin{bmatrix}2m-2\\m-2\end{bmatrix}\equiv(-1)^{m-2}q^{2-\binom{m-1}{2}} \pmod{\Phi_m(q)},$$

and $(-q;q)_{m-1}\equiv 1 \pmod{\Phi_m(q)}$ (see, for example, [4]). The proof of (2b) then follows easily from (8).

3. Proof of Theorems 1.2 and 1.3

Proof of Theorem 1.2. It is easy to see by induction on N that

$$\begin{aligned}&\sum_{k=0}^{N}[4k-1]\frac{(aq^{-1};q^2)_k(q^{-1}/a;q^2)_k(q^{-1};q^2)_k^2}{(aq^2;q^2)_k(q^2/a;q^2)_k(q^2;q^2)_k^2}q^{4k}\\&\quad=\frac{(aq;q^2)_N(q/a;q^2)_N((a+1)^2q^{2N+1}-a(1+q)(1+q^{4N+1}))}{q(a-q)(1-aq)(aq^2;q^2)_N(q^2/a;q^2)_N(-q;q)_N^4}\begin{bmatrix}2N\\N\end{bmatrix}^2. \quad (9)\end{aligned}$$

For $N=(n+1)/2$ or $N=n-1$, we see that $(aq;q^2)_N(q/a;q^2)_N$ contains the factor $(1-aq^n)(1-q^n/a)$. Moreover,

$$\frac{[(n+1)/2]}{[n]}\begin{bmatrix} n \\ (n-1)/2 \end{bmatrix} = \begin{bmatrix} n-1 \\ (n-1)/2 \end{bmatrix}$$

is a polynomial in q. Since $[(n+1)/2]$ and $[n]$ are relatively prime, we conclude that $\left[\begin{smallmatrix} n \\ (n-1)/2 \end{smallmatrix}\right]$ is divisible by $[n]$. Therefore, $\left[\begin{smallmatrix} n+1 \\ (n+1)/2 \end{smallmatrix}\right] = (1+q^{(n+1)/2})\left[\begin{smallmatrix} n \\ (n-1)/2 \end{smallmatrix}\right]$ is also divisible by $[n]$. It is also well known that $\left[\begin{smallmatrix} 2n-2 \\ n-1 \end{smallmatrix}\right]$ is divisible by $[n]$. Moreover, it is easy to see that $[n]$ is relatively prime to $1+q^m$ for any non-negative integer m. The proof then follows from (9) by taking $N=(n+1)/2$ and $N=n-1$. □

Proof of Theorem 1.3. For $a=-1$, the identity (9) reduces to

$$\begin{aligned}\sum_{k=0}^{N}[4k-1]\frac{(q^{-2};q^4)_k^2}{(q^4;q^4)_k^2}q^{4k} &= -\frac{(-q;q^2)_N^2(1+q^{4N+1})}{q(1+q)(-q^2;q^2)_N^2(-q;q)_N^4}\begin{bmatrix} 2N \\ N \end{bmatrix}^2 \\ &= -\frac{(1+q^{4N+1})}{q(1+q)(-q^2;q^2)_N^4}\begin{bmatrix} 2N \\ N \end{bmatrix}_{q^2}^2 \qquad (10)\end{aligned}$$

Note that, in the proof of Theorem 1.2, we have proved that $\left[\begin{smallmatrix} 2N \\ N \end{smallmatrix}\right]_{q^2}$ is divisible by $[n]_{q^2}$ for both $N=(n+1)/2$ and $N=n-1$. Moreover, $[n]_{q^2}$ is relatively prime to $(-q^2;q^2)_m$ for $m\geqslant 0$. Hence the right-hand side of (10) is congruent to 0 modulo $[n]_{q^2}^2$ for $N=(n+1)/2$ or $N=n-1$. To further determine the right-hand side of (10) modulo $[n]_{q^2}^2\Phi_n(q^2)$, we need only to use the same congruences (with $q\mapsto q^2$) used in the proof of Theorem 1.1. □

4. Immediate consequences

Notice that for $n=p^r$ being an odd prime power, $\Phi_{p^r}(q)=[p]_{q^{p^r-1}}$ holds. This observation was used in [15] to extend (1) to a supercongruence modulo $[p^r][p]_{q^{p^r-1}}^3$. In the same vein we immediately deduce from Theorem 1.1 the following result:

Corollary 4.1. *Let p be an odd prime and r a positive integer. Then*

$$\sum_{k=0}^{\frac{p^r+1}{2}}[4k-1]\frac{(q^{-1};q^2)_k^4}{(q^2;q^2)_k^4}q^{4k} \equiv -(1+3q+q^2)[p^r]^4 \pmod{[p^r]^4[p]_{q^{p^r-1}}}, \qquad (11a)$$

and

$$\sum_{k=0}^{p^r-1}[4k-1]\frac{(q^{-1};q^2)_k^4}{(q^2;q^2)_k^4}q^{4k} \equiv -(1+3q+q^2)[p^r]^4 \pmod{[p^r]^4[p]_{q^{p^r-1}}}. \qquad (11b)$$

The $q\to 1$ limiting cases of these two identities yield the following supercongruences:

Corollary 4.2. *Let p be an odd prime and r a positive integer. Then*

$$\sum_{k=0}^{\frac{p^r-1}{2}} \frac{4k+3}{16(k+1)^4\,256^k}\binom{2k}{k}^4 \equiv 1-5p^{4r} \pmod{p^{4r+1}}, \tag{12a}$$

and

$$\sum_{k=0}^{p^r-2} \frac{4k+3}{16(k+1)^4\,256^k}\binom{2k}{k}^4 \equiv 1-5p^{4r} \pmod{p^{4r+1}}. \tag{12b}$$

Similarly, we deduce from Theorem 1.3 the following result:

Corollary 4.3. *Let p be an odd prime and r a positive integer. Then*

$$\sum_{k=0}^{\frac{p^r+1}{2}} [4k-1]\frac{(q^{-2};q^4)_k^2}{(q^4;q^4)_k^2}q^{4k} \equiv -q^{p^r}(1-q+q^2)[p^r]_{q^2}^2 \pmod{[p^r]_{q^2}^2[p]_{q^{2p^r-1}}}, \tag{13a}$$

and

$$\sum_{k=0}^{p^r-1} [4k-1]\frac{(q^{-2};q^4)_k^2}{(q^4;q^4)_k^2}q^{4k} \equiv -(1-q+q^2)[p^r]_{q^2}^2 \pmod{[p^r]_{q^2}^2[p]_{q^{2p^r-1}}}. \tag{13b}$$

The $q\to 1$ limiting cases of these two identities yield the following supercongruences:

Corollary 4.4. *Let p be an odd prime and r a positive integer. Then*

$$\sum_{k=0}^{\frac{p^r-1}{2}} \frac{4k+3}{4(k+1)^2\,16^k}\binom{2k}{k}^2 \equiv 1-p^{2r} \pmod{p^{2r+1}}, \tag{14a}$$

and

$$\sum_{k=0}^{p^r-2} \frac{4k+3}{4(k+1)^2\,16^k}\binom{2k}{k}^2 \equiv 1-p^{2r} \pmod{p^{2r+1}}. \tag{14b}$$

The supercongruences in Corollaries 4.2 and 4.4 are remarkable since they are valid for arbitrarily high prime powers. Swisher [24] had empirically observed several similar but different hypergeometric supercongruences and stated them without proof.

References

[1] O. Gorodetsky, q-Congruences, with applications to supercongruences and the cyclic sieving phenomenon, preprint, May 2018; arXiv:1805.01254.

[2] J. Guillera, WZ pairs and q-analogues of Ramanujan series for $1/\pi$, *J. Difference Equ. Appl.* **24** (2018), 1871–1879.

[3] V.J.W. Guo, A q-analogue of the (J.2) supercongruence of Van Hamme, *J. Math. Anal. Appl.* **466** (2018), 776–788.

[4] V.J.W. Guo, A q-analogue of the (I.2) supercongruence of Van Hamme, *Int. J. Number Theory* **15** (2019), 29–36.

[5] V.J.W. Guo, Proof of a q-congruence conjectured by Tauraso, *Int. J. Number Theory* **15** (2019), 37–41.
[6] V.J.W. Guo, q-Analogues of the (E.2) and (F.2) supercongruences of Van Hamme, *Ramanujan J.* (2018), `https://doi.org/10.1007/s11139-018-0021-z`
[7] V.J.W. Guo, q-Analogues of two "divergent" Ramanujan-type supercongruences, *Ramanujan J.*, `https://doi.org/10.1007/s11139-019-00161-0`
[8] V.J.W. Guo, A q-analogue of a curious supercongruence of Guillera and Zudilin, *J. Difference Equ. Appl.* **25** (2019), 342–350.
[9] V.J.W. Guo, Factors of some truncated basic hypergeometric series, *J. Math. Anal Appl.* **476** (2019), 851–859.
[10] V.J.W. Guo, Some q-congruences with parameters, *Acta Arith.*, to appear.
[11] V.J.W. Guo and H.-X. Ni, A q-congruence involving the Jacobi symbol, *Int. J. Number Theory*, to appear.
[12] V.J.W. Guo and M.J. Schlosser, Some new q-congruences for truncated basic hypergeometric series, *Symmetry* **11**(2) (2019):268.
[13] V.J.W. Guo and M.J. Schlosser, Some q-supercongruences from transformation formulas for basic hypergeometric series, preprint, December 2018; arXiv:1812.06324.
[14] V.J.W. Guo and S.-D. Wang, Factors of sums and alternating sums of products of q-binomial coefficients and powers of q-integers, *Taiwan. J. Math.* 23 (2019), 11–27.
[15] V.J.W. Guo and S.-D. Wang, Some congruences involving fourth powers of central q-binomial coefficients, *Proc. Roy. Soc. Edinburgh Sect. A*, `https://doi.org/10.1017/prm.2018.96`
[16] V.J.W. Guo and W. Zudilin, A q-microscope for supercongruences, *Adv. Math.* **346** (2019), 329–358.
[17] V.J.W. Guo and W. Zudilin, On a q-deformation of modular forms, *J. Math. Anal Appl.* **475** (2019), 1636–1646.
[18] J. Liu, H. Pan, and Y. Zhang, A generalization of Morley's congruence, *Adv. Difference Equ.* (2015) 2015:254.
[19] J.-C. Liu and F. Petrov, Congruences on sums of q-binomial coefficients, preprint, February 2019; arXiv:1902.03851.
[20] H.-X. Ni and H. Pan, Divisibility of some binomial sums, preprint, August 2018; arXiv:1808.03213.
[21] R. Osburn and W. Zudilin, On the (K.2) supercongruence of Van Hamme, *J. Math. Anal. Appl.* 433 (2016), 706–711.
[22] M. Petkovšek, H.S. Wilf, and D. Zeilberger, $A = B$, A K Peters, Ltd., Wellesley, MA, 1996.
[23] A. Straub, Supercongruences for polynomial analogs of the Apéry numbers, *Proc. Amer. Math. Soc.* **147** (2019), 1023–1036.
[24] H. Swisher, On the supercongruence conjectures of van Hamme, *Res. Math. Sci.* (2015) 2:18.
[25] R. Tauraso, q-Analogs of some congruences involving Catalan numbers, *Adv. Appl. Math.* **48** (2009), 603–614.
[26] R. Tauraso, Some q-analogs of congruences for central binomial sums, *Colloq. Math.* **133** (2013), 133–143.
[27] L. Van Hamme, Some conjectures concerning partial sums of generalized hypergeometric series, in: *p-Adic Functional Analysis* (Nijmegen, 1996), Lecture Notes in Pure and Appl. Math. 192, Dekker, New York, 1997, pp. 223–236.
[28] H.S. Wilf and D. Zeilberger, An algorithmic proof theory for hypergeometric (ordinary and "q") multisum/integral identities, *Invent. Math.* **108** (1992), 575–633.
[29] W. Zudilin, Congruences for q-binomial coefficients, *Ann. Comb.*, to appear.

School of Mathematical Sciences, Huaiyin Normal University, Huai'an 223300, Jiangsu, People's Republic of China
E-mail address: jwguo@hytc.edu.cn

Fakultät für Mathematik, Universität Wien, Oskar-Morgenstern-Platz 1, A-1090 Vienna, Austria
E-mail address: michael.schlosser@univie.ac.at