

SIMPLE ROOT SYSTEMS AND PRESENTATIONS FOR CERTAIN COMPLEX REFLECTION GROUPS

JIAN-YI SHI

Department of Mathematics, East China
Normal University, Shanghai, 200062, China
and
Center for Combinatorics, Nankai
University, Tianjin, 300071, China

ABSTRACT. We find all the inequivalent simple root systems for the complex reflection groups G_{12} , G_{24} , G_{25} and G_{26} . Then we give all the non-congruent essential presentations of these groups by generators and relations. The method used in the paper is applicable to any finite (complex) reflection groups.

Introduction.

Shephard and Todd classified all the finite complex reflection groups in paper [5]. Later Cohen gave a more systematic description for these groups in terms of root systems (R, f) and root graphs in [2], in particular, he defined a simple root system (B, w) for the root system of such a group, which is analogous to the corresponding concept for a Coxeter group. In general, for a given finite complex reflection group G , a root system (R, f) is essentially unique but a simple root system (B, w) for (R, f) is not (even up to G -action), e.g., when $G = G_{33}, G_{34}$ (see [1, Table 4]). One can define an equivalence relation on the set of simple root systems for (R, f) (see 1.7). A natural question is to ask

Problem A. *How many inequivalent simple root systems are there in total for any irreducible finite complex reflection group G ?*

Key words and phrases. complex reflection groups, simple root systems, congruence presentations.

Supported by the 973 Project of MST of China, the NSF of China, the SF of the Univ. Doctoral Program of ME of China and the Shanghai Priority Academic Discipline

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

It is well known that any Coxeter group can be presented by generators and relations. A finite complex reflection group G can also be presented in a similar way (see [2]). But such a presentation is not unique for G in general. Different presentations of G may reveal various different properties of G (see [6] for example). Then it is worth to define a congruence relation among the presentations of G (see 1.9) and then to ask

Problem B. *How many non-congruent presentations are there for any irreducible finite complex reflection group G ?*

In the present paper, we solve these problems just for the finite primitive complex reflection groups $G = G_{12}, G_{24}, G_{25}, G_{26}$ (in the notations of [5]). The other finite primitive complex reflection groups except G_{34} will be dealt with subsequently in two papers by my graduate students Li Wang and Peng Zeng (see [8, 9]). We choose to deal with these four groups as they represent four different cases. Among these groups, G_{12} is the only group the number of whose generating reflections is not equal to the dimension of the space it acts. The group G_{24} is generated by three reflections of order 2, G_{25} is generated by three reflections of order 3, while G_{26} is generated by three reflections of different orders: one of order 2 and two of order 3. Our method in dealing with Problem A is based on the knowledge of the action of G on its root system. Then the results so obtained, together with a known presentation of G , will be used in dealing with Problem B. We shall further show that the reduced forms for the presentations of the groups obtained in the paper are all essential (see 7.1 and Theorem 7.3). It is a relatively easy task if one is only content with getting the non-congruent essential presentations for a given finite complex reflection group. But some of such presentations may have very complicated form and hence is not applicable in practice. So in the present paper we are trying to find the forms of the non-congruent essential presentations as simple as possible. Then it becomes quite subtle in finding such forms and even more subtle in proving that they are indeed non-congruent and essential. The methods used in the paper are applicable to any other finite (complex) reflection group.

The contents are organized as follows. Section 1 is served as preliminaries, some definitions and results are collected there. Then we give all the inequivalent simple root systems for the group G_{12} in Section 2, and give all the non-congruent presentations of G_{12} in Section 3. We do the same for the groups G_{24} , G_{25} , G_{26} in Sections 4, 5, 6, respectively. Finally we show in Section 7 that the reduced forms for all the presentations of the groups obtained in the paper are essential.

§1. Preliminaries.

We collect some definitions and results concerning irreducible finite reflection groups, where 1.1-1.5 follow from Cohen's paper [2] except for the definition of a simple root system which follows from [3].

1.1. Let V be a complex vector space of dimension n . A *reflection* on V is a linear transformation on V of finite order with exactly $n - 1$ eigenvalues equal to 1. A *reflection group* G on V is a finite group generated by reflections on V . The group G is *reducible* if it is a direct product of two proper reflection subgroups and *irreducible* otherwise. The action of G on V is said to be *irreducible* if V has no nonzero proper G -invariant subspace. In the present paper we shall always assume that G is irreducible and acts irreducibly on V . A reflection group G on V is called a *real group* or a *Coxeter group* if there is a G -invariant \mathbb{R} -subspace V_0 of V such that the canonical map $\mathbb{C} \otimes_{\mathbb{R}} V_0 \rightarrow V$ is bijective. If this is not the case, G will be called *complex*. (Note that, according to this definition, a real reflection group is not complex.)

Since G is finite, there exists a unitary inner product $(\ , \)$ on V invariant under G . From now on we assume that such an inner product is fixed.

1.2. A *root* of a reflection on V is an eigenvector corresponding to the unique nontrivial eigenvalue of the reflection. A *root of G* is a root of a reflection in G .

Let s be a reflection on V of order $d > 1$. There is a vector $a \in V$ of length 1 and a primitive d -th root ζ of unity such that $s = s_{a,\zeta}$, where $s_{a,\zeta}$ is defined by

$$(1.2.1) \quad s_{a,\zeta}(v) = v + (\zeta - 1)(v, a)a \quad \text{for all } v \in V.$$

We also write $s_{a,d}$ for $s_{a,\zeta}$ if $\zeta = e^{2\pi i/d}$. In this case, we often write $s_{a,d}$ simply by s_a if d is clear from the context.

For each $v \in V$ define $o_G(v)$ to be the order of the (necessarily cyclic) group that consists of the identity and the reflections in G which have v as a root. (This group is $G_W = \{g \in G \mid gu = u \text{ for all } u \in W\}$, where $W = v^\perp$.) Thus $o_G(v) > 1$ if and only if v is a root of G . If a is a root of G , then $o_G(a)$ will be called the *order* of a (with respect to G). We shall denote $o_G(a)$ simply by $o(a)$ when G is clear from the context.

The following results can be shown easily.

Lemma 1.3. (1) $o(gv) = o(v) = o(cv)$ for all $v \in V$, $g \in G$ and $c \in \mathbb{C} \setminus \{0\}$.

(2) $gs_ag^{-1} = s_{g(a)}$ for any $g \in G$ and any root a of G .

1.4. A pair (R, f) is called a *root system* in V , if

- (i) R is a finite set of vectors v spanning V with $|(v, v)| = 1$;
- (ii) $f : R \rightarrow \mathbb{N} \setminus \{1\}$ is a map such that $s_{a, f(a)}R = R$ and $f(s_{a, f(a)}(b)) = f(b)$ for all $a, b \in R$;
- (iii) the group G generated by $\{s_{a, f(a)} \mid a \in R\}$ is a finite reflection group, and for all $a \in R$ and $c \in \mathbb{C}$,

$$ca \in R \iff ca \in Ga.$$

The group G is called the reflection group associated with the root system (R, f) . We have $o_G(a) = f(a)$ for any $a \in R$.

We shall denote a root system (R, f) simply by R when f is clear from the context.

1.5. A *simple root system* is a pair (B, w) , where B is a finite set of vectors spanning V and w is a map from B to $\mathbb{N} \setminus \{1\}$, satisfying the following conditions:

- (i) for all $a, b \in B$, we have $|(a, b)| = 1 \iff a = b$;
- (ii) the group G generated by $S = \{s_{a, w(a)} \mid a \in B\}$ is finite;
- (iii) there is a root system (R, f) with $R = GB$ and $f(a) = w(a)$ for all $a \in B$;
- (iv) the group G cannot be generated by fewer than $|B|$ reflections.

We call the elements of S *simple reflections*. We also call (R, f) the root system of G *generated by B* , and B a *simple root system* for R (or for G).

Note that we do not require B to be linearly independent. If B is linearly independent, then condition (iv) holds automatically.

The above definition of a simple system is considerably weaker than the usual definition for Coxeter groups; in particular, it is not always true that if B_1 and B_2 are simple root systems for the same root system R then there is an element $g \in G$ with $gB_1 = B_2$.

By Lemma 1.3 we see that if $\alpha \in B$ and $\beta \in \mathbb{C}\alpha \cap R$, then $B' = (B \setminus \{\alpha\}) \cup \{\beta\}$ also forms a simple root system for R .

1.6. Let B (resp. B') be a subset of V and $w : B \rightarrow \mathbb{N} \setminus \{1\}$ (resp. $w' : B' \rightarrow \mathbb{N} \setminus \{1\}$) be a function. Let G (resp. G') be the reflection group generated by $\{s_{\alpha, w(\alpha)} \mid \alpha \in B\}$ (resp. $\{s_{\alpha, w'(\alpha)} \mid \alpha \in B'\}$). If there exists a bijection $\phi : B \rightarrow B'$ such that $w(\alpha) = w'(\phi(\alpha))$ and $(\alpha, \beta) = (\phi(\alpha), \phi(\beta))$ for any $\alpha, \beta \in B$. Then by the theory of linear algebra, we see that the assignment $s_{\alpha, w(\alpha)} \mapsto s_{\phi(\alpha), w'(\phi(\alpha))}$ determines a group isomorphism from G to G' .

1.7. Let (R, f) be a root system with G the associated reflection group and B a simple root system for R . Then it is known that as a root system for G , R is determined by G

up to scalar factors (see [3, 1.9]). However, a simple root system B for R is not uniquely determined by G . Two simple root systems (B, w) and (B', w') for (R, f) are *equivalent*, written $B \sim B'$, if there exists a bijection $\phi : B \longrightarrow B'$ such that for any $\alpha, \beta \in B$,

- (1) $w(\alpha) = w'(\phi(\alpha))$ and,
- (2) $\langle s_{\alpha, w(\alpha)}, s_{\beta, w(\beta)} \rangle \cong \langle s_{\phi(\alpha), w'(\phi(\alpha))}, s_{\phi(\beta), w'(\phi(\beta))} \rangle$, where the notation $\langle s, t \rangle$ stands for the group generated by s, t .

In particular, two simple root systems B, B' for R are equivalent if one of the following cases occurs:

- (a) $gB = B'$ for some $g \in G$;
- (b) Condition (2) is replaced by condition (2') below in the above definition of $B \sim B'$:
- (2') $|(\alpha, \beta)| = |(\phi(\alpha), \phi(\beta))|$ for any $\alpha, \beta \in B$.

In general, not all the simple root systems are equivalent for a given reflection group. So it is natural to ask Problem A in Introduction.

It is known that at least one simple root system has been found for any irreducible finite reflection group G (see [7]). We have the following criterion for a subset of a root system to be a simple root system.

Proposition 1.8. *Let G be a finite reflection group with (R, f) the associated root system and B a simple root system for R . Let B' be a subset of R and let $G_{B'}$ be the group generated by the reflections $s_{\alpha, f(\alpha)}$, $\alpha \in B'$.*

- (1) *B' forms a simple root system for G if and only if $G_{B'}B' = R$ and $|B'| = |B|$.*
- (2) *If $B \subseteq G_{B'}B'$ then $G_{B'}B' = R$.*

Proof. (1) The implication “ \implies ” is obvious by the definition of a simple root system. For the reversing implication, we need only check conditions (i) and (ii) in 1.5. Since $G_{B'}B' = R$, we have $G_{B'} = G$. So (ii) follows by the assumption that G is finite. Since (R, f) is a root system, any vector v in R (in particular, when v is in B') satisfies $|(v, v)| = 1$ and hence $|(v, v')| \neq 1$ for any non-proportional $v, v' \in R$. So to show that (B', w') satisfies (i), it is enough to show that B' does not contain two proportional vectors α, β . For otherwise, the reflections with respect to the vectors in the proper subset $B' \setminus \{\beta\}$ of B' would generate the group G , contradicting the assumptions that $|B'| = |B|$ and that B is a simple root system for G .

(2) Any reflection in the group $G_{B'}$ has the form s_{α} with $\alpha \in G_{B'}B'$. Since $B \subseteq G_{B'}B'$ and $G = \langle s_{\beta} \mid \beta \in B \rangle$, we have $G_{B'} \supseteq G$ and hence $G_{B'}$ contains all the reflections s_{β} ,

$\beta \in R$. This implies the inclusion $R \subseteq G_{B'}B'$ and hence $G_{B'}B' = R$. \square

1.9. Given a reflection group G , a *presentation of G by generators and relations* (or a *presentation* in short) is by definition a pair (S, \mathcal{P}) , where

(1) S is a finite generator set for G which consists of reflections, and S has minimal cardinality with this property.

(2) \mathcal{P} is a finite set of relations on S , and any other relation on S is a consequence of the relations in \mathcal{P} .

Two presentations (S, \mathcal{P}) and (S', \mathcal{P}') for G are *congruent*, if there exists a bijection $\eta : S \longrightarrow S'$ such that for any $s, t \in S$,

$$(*) \quad \langle s, t \rangle \cong \langle \eta(s), \eta(t) \rangle.$$

In this case, we see by taking $s = t$ that the order $o(s)$ of s is equal to the order $o(\eta(s))$ of $\eta(s)$ for any $s \in S$.

If there does not exist such a bijection η , then we say that they are *non-congruent*.

For any reflections s, t in G , it is known (see, for example, Koster [4]) that there exists a positive integer k such that $sts\dots = tst\dots$ (k factors on each side). Denote by $n(s, t)$ the smallest such k . It is easily seen that when $o(s) = o(t) = 2$, condition $(*)$ is equivalent to condition

$$(**) \quad n(s, t) = n(\eta(s), \eta(t)).$$

However, when either $o(s), o(t) > 2$ or $\max\{o(s), o(t)\} = 5$, one can show that neither of $(*)$ and $(**)$ implies the other in general by checking all the cases where a complex reflection group is irreducible and is generated by exactly two reflections.

1.10. Given a simple root system (B, w) for G , say $B = \{\alpha_1, \dots, \alpha_r\}$, denote by $\mathbf{n}(B)$ the $\binom{r}{2}$ -tuple $(X_{12}, X_{13}, \dots, X_{1r}, X_{23}, X_{24}, \dots, X_{2r}, \dots, X_{r-1,r})$ with $\binom{r}{2} = \frac{r(r-1)}{2}$, where $X_{ij} = \langle s_{\alpha_i, w(\alpha_i)}, s_{\alpha_j, w(\alpha_j)} \rangle$ for $1 \leq i < j \leq r$. In particular, when w is the constant function 2, we can replace the group X_{ij} by the number $n(s_{\alpha_i, 2}, s_{\alpha_j, 2})$ for any i, j in the definition of $\mathbf{n}(B)$. This is because in that case, X_{ij} becomes a dihedral group which is entirely determined by the number $n(s_{\alpha_i, 2}, s_{\alpha_j, 2})$.

1.11. For each presentation (S, \mathcal{P}) for G , there exists a simple root system (B_S, w_S) for G (called an associated simple root system of (S, \mathcal{P})) such that each element of S has the form $s_{\alpha, w_S(\alpha)}^{k_S(\alpha)}$ for some $\alpha \in B_S$ and some $k_S(\alpha) \in \mathbb{N}$ with $k_S(\alpha)$ coprime to $w_S(\alpha)$. By the minimality for the cardinality of S , we see that for any $\alpha, \beta \in B_S$, $s_{\alpha, w_S(\alpha)}^{k_S(\alpha)} = s_{\beta, w_S(\beta)}^{k_S(\beta)}$ in S if and only if $\alpha = \beta$. By properly choosing generators in a presentation (S, \mathcal{P}) for G ,

we can always make an associated simple root system (B_S, w_S) to satisfy $k_S(\alpha) = 1$ for all $\alpha \in B_S$)

Lemma. *Two presentations of G are congruent if and only if their associated simple root systems are equivalent (see 1.7).*

Proof. It follows directly from the definitions. \square

Remark 1.12. (1) In the present paper, we shall deal with Problems A and B (see Introduction) for the groups $G = G_{12}, G_{24}, G_{25}$ and G_{26} . As a basic work in our method, we calculate the group $\langle s_\alpha, s_\beta \rangle$ (or the value $n(s_\alpha, s_\beta)$ when $f(\alpha) = f(\beta) = 2$) for any pair of reflections $s_\alpha \neq s_\beta$ in G with respect to $\alpha, \beta \in R$, and also calculate all the permutations of R given rise by the action of the reflections $s_\alpha, \alpha \in R$ (these results are not included explicitly in the paper). These can be done by computer in general.

(2) Howlett classified all the non-congruent simple reflection sets for the groups G_{12} and G_{24} by a certain detailed analysis on the group-theoretic structure of these groups in a private communication. The method used in the paper is quite different from Howlett's, and could be applied without essential change for the other finite complex reflection groups.

§2. Simple root systems for the group G_{12} .

2.1. Let ϵ_1, ϵ_2 be an orthonormal basis in a hermitian space V of dimension 2. Let $e_1 = \epsilon_1$ and $e_2 = \frac{-1+\sqrt{2}i}{2}\epsilon_1 + \frac{\sqrt{3}-\sqrt{6}i}{6}\epsilon_2$. Then e_1, e_2 is also a basis of V . Denote by (a, b) a vector $ae_1 + be_2$ in V . Let $R_{12} = \{\pm e_i \mid 1 \leq i \leq 12\}$, where

$$\begin{aligned} e_1 &= (1, 0), & e_2 &= (0, 1), & e_3 &= (1, 1), & e_4 &= (1 - \sqrt{2}i, 1), \\ e_5 &= (-\sqrt{2}i, 1), & e_6 &= (1, 1 + \sqrt{2}i), & e_7 &= (1, \sqrt{2}i), & e_8 &= (\sqrt{2}i, -1 + \sqrt{2}i), \\ e_9 &= (1 + \sqrt{2}i, \sqrt{2}i), & e_{10} &= (1 + \sqrt{2}i, -1 + \sqrt{2}i), & e_{11} &= (1 - \sqrt{2}i, 2), & e_{12} &= (2, 1 + \sqrt{2}i). \end{aligned}$$

Let s_i be the reflection in V with respect to e_i for $1 \leq i \leq 12$. Then the action of s_i on e_j , $1 \leq i, j \leq 3$, are as follows.

e_i	$s_1(e_i)$	$s_2(e_i)$	$s_3(e_i)$
e_1	$-e_1$	e_6	e_8
e_2	e_4	$-e_2$	$-e_9$
e_3	e_5	e_7	$-e_3$

The group generated by the set $S_1 = \{s_1, s_2, s_3\}$ is G_{12} according to the notation by Shephard-Todd (see [5] and [7]). R_{12} is a root system for G_{12} with $B_1 = \{e_1, e_2, e_3\}$ a simple root system by [1] or [7].

2.2. By Proposition 1.8, we see that any simple root system B for the group G_{12} contains exactly three vectors. Since R_{12} is transitive under G_{12} , we may assume $e_1 \in B$ up to G_{12} -action. Consider all the possible orders $o(s_1 s_i)$ of the product $s_1 s_i$ for $1 < i \leq 12$. Then $o(s_1 s_{11})=2$, $o(s_1 s_i)$ is 3 for $i \in \{6, 8, 10, 12\}$, 4 for $i \in \{7, 9\}$ and 6 for $i \in \{2, 3, 4, 5\}$. Let G_B be the group generated by the reflections with respect to the vectors in B . It is easily checked that no $B \subset R_{12}$ with $e_1, e_{11} \in B$ and $|B| = 3$ satisfies the equation $G_B B = R_{12}$. So no simple root system for G_{12} contains a pair of orthogonal roots. We can also show that any $B \in \{\{e_1, e_6, e_{10}\}, \{e_1, e_8, e_{12}\}\}$ satisfies the condition $G_B B \subsetneq R_{12}$. So in a simple root system there must exist some pair of roots α, β with $n(s_\alpha, s_\beta) > 3$ (see 1.9).

2.4. We can show that any of the following $B_i \subset R_{12}$ satisfies the equation $G_{B_i} B_i = R_{12}$ and hence is a simple root system for G_{12} by Proposition 1.8.

$$B_1 = \{e_1, e_2, e_3\}, \quad B_2 = \{e_1, e_2, e_{10}\}, \quad B_3 = \{e_1, e_2, e_7\}, \quad B_4 = \{e_1, e_2, e_8\}, \quad B_5 = \{e_1, e_3, e_8\}.$$

Recall the notation $\mathbf{n}(B)$ in 1.10. We have $\mathbf{n}(B_1) = (6, 6, 6)$, $\mathbf{n}(B_2) = (6, 3, 3)$, $\mathbf{n}(B_3) = (6, 3, 4)$, $\mathbf{n}(B_4) = (6, 4, 6)$ and $\mathbf{n}(B_5) = (4, 3, 3)$. This implies that B_i , $1 \leq i \leq 5$, are five inequivalent simple root systems for G_{12} . By 1.6 and by the knowledge of the values $n(s_\alpha, s_\beta)$ for all the non-proportional pairs α, β in R_{12} , we can further show that there is no more inequivalent simple root system for G_{12} at all.

§3. Presentations of the group G_{12} .

In this section, we shall find all the non-congruent presentations of the complex reflection group G_{12} by generators and relations according to the results in 2.4 (see Propositions 3.2-3.6).

3.1. Suppose that we are given a complex reflection group G , the associated root system R of G , and all the inequivalent simple root systems B_k , $1 \leq k \leq r$, for R . The strategy for finding all the non-congruent presentations of G is as follows. We start with the presentation of G given by Shephard-Todd which corresponds to a simple root system, say B_1 , for R . Let S_k be the reflection set associated to the simple root system B_k . We make use of the relations among the simple root systems B_k , $1 \leq k \leq r$, to establish some transition

among the S_k 's by Lemma 1.3. Then for each $1 < k \leq r$, we get a presentation of G with the generator set S_k by some related transition. Suppose that we are given a presentation (S_h, \mathcal{P}) of G with the generator set S_h and the transition between the sets S_h and S_k . Then the simplest method to produce a presentation of G with the generator set S_k is just to substitute the reflections of S_h by those of S_k in the relations of \mathcal{P} by the related transition. But it may happen that sometimes the resulting relations we get in the new presentation of G are quite complicated and hence are not very useful in practice. So we shall not always use such a method.

3.2. It is well known that the group G_{12} is presented by the generator set $S = \{s, u, t\}$ and the relations (see [1]):

$$(1) \ s^2 = u^2 = t^2 = 1.$$

$$(2) \ suts = utsu = tsut.$$

The reflections s_1, s_2 and s_3 in V satisfy relations (1), (2) with s_1, s_2, s_3 in the places of s, u, t respectively. Since $B_1 = \{e_1, e_2, e_3\}$ is a simple root system for G_{12} , the assignment $\rho(s) = s_1, \rho(u) = s_2$ and $\rho(t) = s_3$ determines a faithful representation ρ of the group G_{12} to $\text{GL}(V)$. This implies part (1) of the following

Proposition 3.3. (1) *The group G_{12} can be presented by the generator set $S_1 = \{s_1, s_2, s_3\}$ and the relations:*

$$(a1) \ s_1^2 = s_2^2 = s_3^2 = 1.$$

$$(a2) \ s_1 s_2 s_3 s_1 = s_2 s_3 s_1 s_2.$$

$$(a3) \ s_1 s_2 s_3 s_1 = s_3 s_1 s_2 s_3.$$

(2) *Under the presentation of G_{12} in (1), we have*

$$(a4) \ s_1 s_2 s_1 s_2 s_1 s_2 = s_2 s_1 s_2 s_1 s_2 s_1.$$

$$(a5) \ s_2 s_3 s_2 s_3 s_2 s_3 = s_3 s_2 s_3 s_2 s_3 s_2.$$

$$(a6) \ s_1 s_3 s_1 s_3 s_1 s_3 = s_3 s_1 s_3 s_1 s_3 s_1.$$

$$(a7) \ o(s_1 s_2 s_3) = 8.$$

Proof. We need only show (2). Let $x = s_1 s_2 s_3 s_1$ and $y = s_1 s_2 s_3$. Then $x^6 = y^8$ and $o(x) = o(s_1 s_2) = o(s_1 s_3) = o(s_2 s_3)$ by (a1)-(a3). Thus to show (a4)-(a7), it suffices to show that $x^6 = 1, x^2 \neq 1$ and $x^3 \neq 1$, where 1 is the identity transformation in V . These can be shown by observing the action of x^6, x^2, x^3 on the basis elements e_1, e_2 in V . \square

3.4. We have, by Lemma 1.3 (2), that

$$(3.4.1) \quad s_{10} = s_2 s_3 s_2 s_3 s_2 \quad \text{and} \quad s_3 = s_1 s_2 s_{10} s_1 s_{10} s_2 s_1.$$

Proposition. (1) *The group G_{12} can be presented by the generator set $S_2 = \{s_1, s_2, s_{10}\}$ and the relations:*

$$(b1) \quad s_1^2 = s_2^2 = s_{10}^2 = 1.$$

$$(b2) \quad s_1 s_2 s_1 s_2 s_1 s_2 = s_2 s_1 s_2 s_1 s_2 s_1.$$

$$(b3) \quad s_1 s_{10} s_1 = s_{10} s_1 s_{10}.$$

$$(b4) \quad s_1 s_{10} s_2 \cdot s_1 = s_2 \cdot s_1 s_{10} s_2.$$

(2) *Under the presentation of G_{12} in (1), we have*

$$(b5) \quad s_2 s_{10} s_2 = s_{10} s_2 s_{10}.$$

Proof. (b5) follows from (b1), (b3) and (b4). This shows (2) under the assumption of (1). Now we show (1). We must show that relations (a1)-(a3) in Proposition 3.3 are equivalent to (b1)-(b4) here under transition (3.4.1). Clearly, (a1) is equivalent to (b1). So it remains to show that relations (a2)-(a3) are equivalent to (b2)-(b4) under the assumption of (3.4.1) and (a1) (hence (b1)).

(i) First assume (a2)-(a3). Hence (a4)-(a7) hold by Proposition 3.3 (2). Now (b2) is the same as (a4). By (a2)-(a3), we have

$$(3.4.2) \quad s_3 s_2 s_3 s_1 = s_1 s_2 s_3 s_2.$$

(b4) is amount to

$$(3.4.3) \quad s_1 \cdot s_2 s_3 s_2 s_3 s_2 \cdot s_2 s_1 s_2 \cdot s_2 s_3 s_2 s_3 s_2 \cdot s_1 s_2 = 1$$

By (3.4.2) and (a1)-(a7), LHS of (3.4.3) is equal to

$$\begin{aligned} & s_1 s_2 s_3 s_2 s_3 s_1 s_3 s_2 s_3 s_2 s_1 s_2 = s_1 s_2 s_1 s_2 s_3 s_2 s_3 s_2 s_3 s_2 s_1 s_2 \\ & = s_1 s_2 s_1 \cdot s_3 s_2 s_3 s_2 s_3 \cdot s_1 s_2 = s_1 s_2 s_1 s_3 s_2 s_1 s_2 s_3 = s_1 s_2 \cdot s_2 s_1 s_3 s_2 \cdot s_2 s_3 = 1. \end{aligned}$$

So (3.4.3) follows and hence we get (b4).

For (b3), we need show

$$(3.4.4) \quad s_1 \cdot s_2 s_3 s_2 s_3 s_2 \cdot s_1 \cdot s_2 s_3 s_2 s_3 s_2 \cdot s_1 \cdot s_2 s_3 s_2 s_3 s_2 = 1.$$

By (3.4.2) and (a1)-(a7), LHS of (3.4.4) becomes

$$\begin{aligned} & s_1 s_2 s_3 \cdot s_1 s_3 s_2 s_3 \cdot s_2 s_3 s_2 s_3 s_2 \cdot s_3 s_2 s_3 s_1 \cdot s_3 s_2 \\ &= s_1 s_2 s_3 s_1 \cdot s_2 s_1 s_3 s_2 = s_1 s_2 s_3 s_1 \cdot s_1 s_3 s_2 s_1 = 1. \end{aligned}$$

This shows (3.4.4) and hence (b3) follows.

(ii) Next assume (b2)-(b4). Hence (b5) holds also. To show (a2)-(a3), we must show the following two equations:

$$(3.4.5) \quad s_1 s_2 \cdot s_1 s_2 s_{10} s_1 s_{10} s_2 s_1 \cdot s_1 = s_2 \cdot s_1 s_2 s_{10} s_1 s_{10} s_2 s_1 \cdot s_1 s_2$$

$$(3.4.6) \quad s_1 s_2 \cdot s_1 s_2 s_{10} s_1 s_{10} s_2 s_1 \cdot s_1 = s_1 s_2 s_{10} s_1 s_{10} s_2 s_1 \cdot s_1 s_2 \cdot s_1 s_2 s_{10} s_1 s_{10} s_2 s_1.$$

Now by (b1)-(b5), we have

$$\begin{aligned} (3.4.5) &\iff s_1 s_2 \cdot s_1 s_2 s_{10} s_1 s_{10} s_2 \cdot s_{10} s_1 s_{10} s_2 s_1 \cdot s_2 = 1 \\ &\iff s_1 s_2 s_1 s_2 \cdot s_1 s_{10} s_1 s_2 s_1 s_{10} s_1 \cdot s_2 s_1 s_2 = 1 \\ &\iff s_1 s_2 s_1 s_2 s_1 s_2 s_1 s_2 \cdot s_1 s_2 s_1 s_2 = 1. \end{aligned}$$

The last equation follows by (b2) and hence we get (3.4.5).

Finally, by (b1)-(b5), we have

$$\begin{aligned} (3.4.6) &\iff s_1 s_2 s_{10} s_1 s_{10} s_2 \cdot s_1 s_2 s_{10} s_1 s_{10} s_2 s_1 \cdot s_{10} s_1 s_{10} = 1 \\ &\iff s_1 s_2 s_{10} s_1 \cdot s_1 s_2 s_1 \cdot s_1 s_{10} s_2 s_1 \cdot s_1 s_{10} s_1 = 1 \\ &\iff s_2 s_{10} s_2 s_{10} s_2 s_{10} = 1. \end{aligned}$$

The last equation follows by (b5) and hence (3.4.6) is shown. \square

3.5. We have

$$(3.5.1) \quad s_7 = s_2 s_3 s_2 \quad \text{and} \quad s_3 = s_2 s_7 s_2.$$

Proposition. (1) The group G_{12} can be presented by the generator set $S_3 = \{s_1, s_2, s_7\}$ and the relations:

$$(c1) \quad s_1^2 = s_2^2 = s_7^2 = 1.$$

$$(c2) \ s_1 s_2 s_7 \cdot s_1 = s_2 \cdot s_1 s_2 s_7.$$

$$(c3) \ s_7 s_2 s_1 \cdot s_7 = s_2 \cdot s_7 s_2 s_1.$$

(2) Under the presentation of G_{12} in (1), we have

$$(c4) \ s_1 s_7 s_1 s_7 = s_7 s_1 s_7 s_1.$$

$$(c5) \ s_1 s_2 s_1 s_2 s_1 s_2 = s_2 s_1 s_2 s_1 s_2 s_1.$$

$$(c6) \ s_2 s_7 s_2 s_7 s_2 s_7 = s_7 s_2 s_7 s_2 s_7 s_2.$$

Proof. (1) We must show that relations (a1)-(a3) in Proposition 3.3 are equivalent to (c1)-(c3) here under transition (3.5.1).

By (3.5.1), relation (a1) is equivalent to (c1). Under the assumption of (3.5.1) and (a1) (hence (c1)), we have

$$(a2) \iff s_1 s_2 \cdot s_2 s_7 s_2 \cdot s_1 = s_2 \cdot s_2 s_7 s_2 \cdot s_1 s_2$$

$$\iff s_1 s_7 s_2 s_1 = s_7 s_2 s_1 s_2 \iff (c2)$$

□

Then under the assumption of relations (a1)-(a2) (hence (c1)-(c2)), we can show that relation (a3) is equivalent to (c3).

(c5) is the same as (a4). So (c5) follows from (c1)-(c3) by the equivalence between (a1)-(a3) and (c1)-(c3). Then one can easily deduce the equations

$$(3.5.1) \quad s_1 s_7 s_1 s_7 = s_1 s_2 s_1 s_2 s_1 s_2 = s_2 s_7 s_2 s_7 s_2 s_7.$$

from (c1)-(c3). So (c4) and (c6) follow from (c5) and (c1). □

3.6. We have

$$(3.6.1) \quad s_8 = s_3 s_1 s_3, \quad \text{and} \quad s_3 = s_1 s_2 s_8 s_2 s_1.$$

Proposition. (1) The group G_{12} can be presented by the generator set $S_4 = \{s_1, s_2, s_8\}$ and the relations:

$$(d1) \ s_1^2 = s_2^2 = s_8^2 = 1.$$

$$(d2) \ s_2 s_1 s_2 s_8 \cdot s_2 = s_1 \cdot s_2 s_1 s_2 s_8.$$

$$(d3) \ s_8 \cdot s_1 s_2 s_8 s_2 = s_1 s_2 s_8 s_2 \cdot s_1.$$

(2) Under the presentation (1) of G_{12} , we have

$$(d4) \ s_1 s_8 s_1 = s_8 s_1 s_8.$$

$$(d5) \ s_2 s_8 s_2 s_8 = s_8 s_2 s_8 s_2.$$

$$(d6) \ s_1 s_2 s_1 s_2 s_1 s_2 = s_2 s_1 s_2 s_1 s_2 s_1.$$

Proof. For (1), we must show that relations (a1)-(a3) in Proposition 3.3 are equivalent to (d1)-(d3) here. (a1) is equivalent to (d1) by (3.6.1). Then under relations (3.6.1) and (a1) (hence (d1)), we have

$$(a2) \iff s_1 s_2 s_1 s_2 s_8 s_2 = s_2 s_1 s_2 s_8 \iff (d2)$$

$$(a3) \iff s_1 s_2 s_1 s_2 s_8 s_2 = s_1 s_2 s_8 s_1 s_2 s_8 s_2 s_1 \iff (d3).$$

This shows (1). Now we show (d4)-(d6) under the presentation (1) of G_{12} . (d6) is the same as (a4). Rewrite (d2) as

$$(3.6.2) \quad s_1 s_2 s_1 s_2 s_1 s_2 = s_2 s_8 s_2 s_8$$

by (d1), this implies (d5)-(d6) from (d1)-(d3) by the equivalence between (a1)-(a3) and (d1)-(d3). Rewrite (d3) as

$$(3.6.3) \quad s_1 s_8 s_1 = s_2 s_8 s_2 s_1 s_2 s_8 s_2.$$

Then to show (d4), it suffices to show that

$$(3.6.4) \quad s_8 s_1 s_8 = s_2 s_8 s_2 s_1 s_2 s_8 s_2.$$

That is to show

$$(3.6.5) \quad s_1 \cdot s_2 s_8 s_2 s_8 = s_2 s_8 s_2 s_8 \cdot s_1,$$

which follows by (3.6.2) and (d6). Hence (d4) is shown. \square

3.7. We have

$$(3.7.1) \quad s_8 = s_1 s_7 s_2 s_7 s_1 \quad \text{and} \quad s_2 = s_7 s_1 s_8 s_1 s_7.$$

Proposition. *The group G_{12} can be presented by the generator set $S_5 = \{s_1, s_7, s_8\}$ and the relations:*

$$(e1) \ s_1^2 = s_7^2 = s_8^2 = 1.$$

$$(e2) \ s_1 s_7 s_1 s_7 = s_7 s_1 s_7 s_1.$$

$$(e3) \ s_1 s_8 s_1 = s_8 s_1 s_8.$$

$$(e4) \ s_7 s_8 s_7 = s_8 s_7 s_8.$$

$$(e5) \ s_1 s_7 s_1 s_7 \cdot s_8 = s_8 \cdot s_1 s_7 s_1 s_7.$$

Proof. We must show that relations (c1)-(c3) are equivalent to (e1)-(e5) under transition (3.7.1). Clearly, (c1) is equivalent to (e1). It remains to show that relations (c2)-(c3) are equivalent to (e2)-(e5) under the assumption of (3.7.1) and (c1) (hence (e1)).

(i) First assume (c2)-(c3). Hence (c4)-(c6) hold by Proposition 3.5 (2). (e2) is the same as (c4). (e3) is equivalent to

$$(3.7.2) \quad s_1 \cdot s_1 s_7 s_2 s_7 s_1 \cdot s_1 \cdot s_1 s_7 s_2 s_7 s_1 \cdot s_1 \cdot s_1 s_7 s_2 s_7 s_1 = 1$$

By (c1)-(c4), LHS of (3.7.2) is equal to

$$\begin{aligned} & s_7 s_2 s_7 s_1 \cdot s_7 s_2 s_7 s_1 \cdot s_7 s_2 s_7 s_1 = s_7 s_2 s_1 \cdot s_7 s_1 s_7 s_1 \cdot s_2 s_7 s_1 \cdot s_7 s_2 s_7 s_1 \\ & = s_2 s_7 s_2 s_1 \cdot s_1 \cdot s_1 s_2 s_7 s_2 s_1 \cdot s_7 s_2 s_7 s_1 = s_2 s_7 s_2 s_1 \cdot s_7 s_2 s_1 \cdot s_2 s_7 s_1 \\ & = s_7 s_2 s_1 \cdot s_1 s_2 s_7 = 1. \end{aligned}$$

This implies (e3). (e4) can be shown similarly. Finally, (e5) follows from (c6) by substitution of $s_2 = s_7 s_1 s_8 s_1 s_7$.

(ii) Next assume (e2)-(e5). By (3.7.1), we get from (e5) that

$$s_1 s_7 s_1 s_7 \cdot s_1 s_7 s_2 s_7 s_1 = s_1 s_7 s_2 s_7 s_1 \cdot s_1 s_7 s_1 s_7$$

and hence

$$(3.7.3) \quad s_1 s_7 s_1 s_7 \cdot s_2 = s_2 \cdot s_1 s_7 s_1 s_7.$$

By (3.7.1) and (3.7.3), we get

$$\begin{aligned} (e3) & \implies s_1 \cdot s_7 s_2 s_7 \cdot s_1 = s_7 s_2 s_7 \cdot s_1 \cdot s_7 s_2 s_7 \\ & \implies s_1 s_7 s_1 s_7 \cdot s_2 \cdot s_7 s_1 s_7 = s_1 s_2 \cdot s_7 s_1 s_7 \cdot s_2 \\ & \implies s_2 \cdot s_1 s_7 s_1 s_7 \cdot s_7 s_1 s_7 = s_1 s_2 \cdot s_7 s_1 s_7 \cdot s_2 \\ & \implies s_1 \cdot s_7 s_2 s_1 = s_7 s_2 s_1 \cdot s_2 \\ & \implies (c2). \end{aligned}$$

Similarly, (c3) follows from (e4) by (3.7.1) and (3.7.3). \square

§4. Simple root systems and presentations for the group G_{24} .

In this section, we shall find all the inequivalent simple root systems for the complex reflection group G_{24} (see 4.2). Then we shall give all the non-congruent presentations of the group G_{24} by generators and relations according to these inequivalent simple root systems (see Propositions 4.3-4.5).

4.1. Let V be a hermitian space of dimension 3 with an orthonormal basis $\epsilon_1, \epsilon_2, \epsilon_3$. Then $e_1 = \epsilon_1, e_2 = \frac{1}{2}(-\epsilon_1 + \epsilon_2 + \alpha\epsilon_3)$ and $e_3 = \frac{\bar{\alpha}}{2}(\epsilon_2 + \epsilon_3)$ also form a basis of V , where $\bar{\alpha}$ denotes the complex conjugate of $\alpha = \frac{1}{2}(1 + i\sqrt{7})$, the latter is a root of the equation $x^2 - x + 2 = 0$. Denote by (a, b, c) a vector $ae_1 + be_2 + ce_3$. Let $R_{24} = \{\pm e_i \mid 1 \leq i \leq 21\}$, where

$$\begin{aligned}
 e_1 &= (1, 0, 0) & e_2 &= (0, 1, 0) & e_3 &= (0, 0, 1) & e_4 &= (-1, 1, 0) \\
 e_5 &= (-\bar{\alpha}, 0, 1) & e_6 &= (0, \alpha, 1) & e_7 &= (1, 0, -\alpha) & e_8 &= (0, 1, \bar{\alpha}) \\
 e_9 &= (-1, 1, 1) & e_{10} &= (-\bar{\alpha}, 1, 1) & e_{11} &= (-1, \alpha, 1) & e_{12} &= (1, \bar{\alpha}, -\alpha) \\
 e_{13} &= (\alpha, 1, \bar{\alpha}) & e_{14} &= (-\bar{\alpha}, 1, 1 + \bar{\alpha}) & e_{15} &= (-1, \alpha, 1 + \alpha) & e_{16} &= (\alpha, \bar{\alpha}, -\alpha) \\
 e_{17} &= (1, \bar{\alpha}, -1 - \alpha) & e_{18} &= (\alpha, 1, 1 + \bar{\alpha}) & e_{19} &= (-\bar{\alpha}, 2, 1 + \bar{\alpha}) & e_{20} &= (-2, \alpha, 1 + \alpha) \\
 e_{21} &= (\alpha, \bar{\alpha}, \bar{\alpha}).
 \end{aligned}$$

Denote by s_i the reflection in V with respect to the vector e_i , $1 \leq i \leq 21$. The action of s_i on e_j , $i, j = 1, 2, 3$, are as follows.

e_i	$s_1(e_i)$	$s_2(e_i)$	$s_3(e_i)$
e_1	$-e_1$	$-e_4$	e_7
e_2	e_4	$-e_2$	e_8
e_3	e_5	e_6	$-e_3$

The group generated by s_1, s_2, s_3 is G_{24} . R_{24} is the root system for G_{24} with $B_1 = \{e_1, e_2, e_3\}$ a simple root system (see [7]).

4.2. We want to find all the inequivalent simple root systems B_k , $k \geq 1$, for G_{24} . By Proposition 1.8, we have $|B_k| = 3$. Since R_{24} is transitive under G_{24} , we may assume $e_1 \in B_k$ up to G_{24} -action. The number $n(s_1, s_i)$ (see 1.9) is 2 for $i \in \{7, 17, 19, 21\}$, 3 for $i \in \{2, 4, 6, 11, 14, 15, 18, 20\}$ and 4 for $i \in \{3, 5, 8, 9, 10, 12, 13, 16\}$.

Any finite reflection group generated by s_1, s_i, s_j with $i \in \{7, 17, 19, 21\}$ is a homomorphic image of the Coxeter group of type A_3 , B_3 or \tilde{C}_2 , hence it has order ≤ 48 and is not

isomorphic to G_{24} (G_{24} has order 336). So a simple root system B_k for G_{24} contains no pair of orthogonal roots. By 1.6, Proposition 1.8 and the knowledge of the values $n(s_i, s_j)$, $1 \leq i, j \leq 21$, we see that the following are all the inequivalent simple root systems for R_{24} :

$$B_1 = \{e_1, e_2, e_3\}, \quad B_2 = \{e_1, e_3, e_4\} \quad \text{and} \quad B_3 = \{e_1, e_3, e_8\}.$$

We have $\mathbf{n}(B_1) = (3, 4, 4)$, $\mathbf{n}(B_2) = (4, 3, 3)$ and $\mathbf{n}(B_3) = (4, 4, 4)$ (see 1.10).

4.3. Next we shall find all the non-congruent presentations of G_{24} by generators and relations.

Proposition. *The group G_{24} can be presented by the generator set $S_1 = \{s_1, s_2, s_3\}$ and the relations:*

- (a1) $s_1^2 = s_2^2 = s_3^2 = 1$.
- (a2) $s_1 s_3 s_1 s_3 = s_3 s_1 s_3 s_1$.
- (a3) $s_1 s_2 s_1 = s_2 s_1 s_2$.
- (a4) $s_2 s_3 s_2 s_3 = s_3 s_2 s_3 s_2$.
- (a5) $s_3 \cdot s_2 s_1 s_2 s_3 s_2 = s_2 s_1 s_2 s_3 s_2 \cdot s_1$.

Proof. This follows by [1, Table 4] and by the argument as for Proposition 3.3 (1). \square

4.4. We have

$$(4.4.1) \quad s_4 = s_1 s_2 s_1 \quad \text{and} \quad s_2 = s_1 s_4 s_1.$$

Proposition. *The group G_{24} can be presented by the generator set $S_2 = \{s_1, s_3, s_4\}$ and the relations:*

- (b1) $s_1^2 = s_3^2 = s_4^2 = 1$.
- (b2) $s_1 s_3 s_1 s_3 = s_3 s_1 s_3 s_1$.
- (b3) $s_1 s_4 s_1 = s_4 s_1 s_4$.
- (b4) $s_3 s_4 s_3 = s_4 s_3 s_4$.
- (b5) $s_1 s_3 s_1 \cdot s_4 s_1 s_3 s_1 s_4 = s_4 s_1 s_3 s_1 s_4 \cdot s_1 s_3 s_1$.

Proof. Under transition (4.4.1), relation (a1) is equivalent to (b1). Also, (a2) is the same as (b2). Then under relations (4.4.1) and (a1) (hence (b1)), relation (a3) (resp. (a4)) is equivalent to (b3) (resp. (b5)). Finally, under relations (4.4.1), (a1), (a3), (b1) and (b3), relation (a5) is equivalent to (b4). So our result follows by Proposition 4.3. \square

4.5. We have

$$(4.5.1) \quad s_8 = s_3 s_2 s_3 \quad \text{and} \quad s_2 = s_3 s_8 s_3.$$

Proposition. (1) *The group G_{24} can be presented by the generator set $S_3 = \{s_1, s_3, s_8\}$ and the relations:*

$$(c1) \quad s_1^2 = s_3^2 = s_8^2 = 1.$$

$$(c2) \quad s_1 s_3 s_1 s_3 = s_3 s_1 s_3 s_1.$$

$$(c3) \quad s_3 s_8 s_3 s_8 = s_8 s_3 s_8 s_3.$$

$$(c4) \quad s_3 s_1 s_3 s_8 s_3 \cdot s_1 = s_8 \cdot s_3 s_1 s_3 s_8 s_3.$$

$$(c5) \quad s_8 s_1 s_8 s_3 s_8 \cdot s_1 = s_3 \cdot s_8 s_1 s_8 s_3 s_8.$$

(2) *Under the presentation of G_{24} in (1), we have*

$$(c6) \quad s_1 s_8 s_1 s_8 = s_8 s_1 s_8 s_1.$$

Proof. Under transition (4.5.1), relation (a1) is equivalent to (c1). Also, (a2) is the same as (c2). Then under relations (4.5.1) and (a1) (hence (c1)), relation (a4) (resp. (a3)) is equivalent to (c3) (resp. (c4)). Then under relations (a1) and (a4) (hence (c1) and (c3)), relation (a5) is equivalent to (c5). This implies (1). Now we show (2). Right-multiplying both sides of (c4) by the inverse of the corresponding sides of (c5) and then applying (c3), we get

$$s_8 s_1 \cdot s_3 s_8 \cdot s_1 s_3 = s_3 s_8 s_1 \cdot s_3 s_8 \cdot s_1 s_3 s_8$$

and hence

$$(4.5.2) \quad s_3 s_8 s_1 s_3 s_8 \cdot s_1 s_3 = s_8 s_1 \cdot s_3 s_8 s_1 s_3 s_8.$$

So for any $k \in \mathbb{N}$, we have

$$s_3 s_8 s_1 s_3 s_8 \cdot (s_1 s_3)^k = (s_8 s_1)^k \cdot s_3 s_8 s_1 s_3 s_8.$$

This implies that $(s_1 s_3)^k = 1$ if and only if $(s_8 s_1)^k = 1$. So (c6) follows by (c2). \square

§5. Simple root systems and presentations for the group G_{25} .

In this section, we deal with the complex reflection group G_{25} . Comparing with the groups G_{12} , G_{24} , a notable difference of G_{25} is that all of its reflections have order 3, rather than 2. There are two inequivalent simple root systems and two non-congruent presentations

for G_{25} (see 5.2 and Propositions 5.3, 5.4). The presentation given in Corollary 5.4 is congruent to that in Proposition 5.4.

5.1. Let V, ϵ_i be as in 4.1. Let $e_1 = \bar{\omega}\epsilon_3, e_2 = 3^{-1/2}i(\epsilon_1 + \epsilon_2 + \epsilon_3)$ and $e_3 = \bar{\omega}\epsilon_2$, where ω is a primitive cubic root of unity and $\bar{\omega}$ its complex conjugate. Then $B_1 = \{e_1, e_2, e_3\}$ forms a basis of V . Denote by (a, b, c) a vector $ae_1 + be_2 + ce_3$. Let $R_{25} = \{(-\omega)^k e_i \mid 1 \leq i \leq 12, 0 \leq k < 6\}$, where

$$\begin{aligned} e_1 &= (1, 0, 0), & e_2 &= (0, 1, 0), & e_3 &= (0, 0, 1), & e_4 &= (-\bar{\omega}, 1, 0), \\ e_5 &= (1, 1, 0), & e_6 &= (0, 1, 1), & e_7 &= (0, 1, -\bar{\omega}), & e_8 &= (-\bar{\omega}, 1, 1), \\ e_9 &= (1, 1, 1), & e_{10} &= (1, 1, -\bar{\omega}), & e_{11} &= (-\bar{\omega}, 1, -\bar{\omega}), & e_{12} &= (-\bar{\omega}, 1 - \bar{\omega}, -\bar{\omega}). \end{aligned}$$

Let s_i be the reflection in V with respect to $e_i, 1 \leq i \leq 12$. Then the group generated by $S_1 = \{s_1, s_2, s_3\}$ is the complex reflection group G_{25} (in the notation of [5]). The set R_{25} forms a root system of G_{25} with B_1 a simple root system (see [7]). The action of S_1 on B_1 is as follows.

e_i	$s_1(e_i)$	$s_2(e_i)$	$s_3(e_i)$
e_1	ωe_1	$-\omega e_4$	e_1
e_2	e_5	ωe_2	e_6
e_3	e_3	$-\omega e_7$	ωe_3

5.2. We want to find all the inequivalent simple root systems $B_k, k \geq 1$, for G_{25} . By Proposition 1.8, we have $|B_k| = 3$. Since R_{25} is transitive under G_{25} , we may assume $e_1 \in B_k$ up to G_{25} -action. Let H_{ij} be the subgroup of G_{25} generated by s_i, s_j for $1 \leq i \neq j \leq 12$. Then by the action on R_{25} of the subgroups H_{1i} for any $1 < i \leq 12$, we see that H_{1i} is isomorphic to the elementary 3-group \mathbb{Z}_3^2 for $i = 3, 12$, and is isomorphic to the group G_4 (in the notation of [5]) for all the other i . From this fact, it is easily seen that all the inequivalent simple root systems for R_{25} are as follows.

$$B_1 = \{e_1, e_2, e_3\}, B_2 = \{e_1, e_2, e_6\}.$$

We have $\mathbf{n}(B_1) = (G_4, G_4, \mathbb{Z}_3^2)$ and $\mathbf{n}(B_2) = (G_4, G_4, G_4)$ (see 1.10).

5.3. Next we shall find all the non-congruent presentations of G_{25} by generators and relations according to the result in 5.2.

Proposition. *The group G_{25} can be presented by the generator set $S_1 = \{s_1, s_2, s_3\}$ and the relations:*

$$(a1) \ s_1^3 = s_2^3 = s_3^3 = 1;$$

$$(a2) \ s_1 s_2 s_1 = s_2 s_1 s_2;$$

$$(a3) \ s_2 s_3 s_2 = s_3 s_2 s_3;$$

$$(a4) \ s_1 s_3 = s_3 s_1.$$

Proof. This follows by [1, Table 1]. \square

5.4. We have

$$(5.4.1) \quad s_6 = s_2^{-1} s_3 s_2 \quad \text{and} \quad s_3 = s_2 s_6 s_2^{-1}.$$

Proposition. (1) The group G_{25} can be presented by the generator set $S_2 = \{s_1, s_2, s_6\}$ and the relations:

$$(b1) \ s_1^3 = s_2^3 = s_6^3 = 1;$$

$$(b2) \ s_1 s_2 s_1 = s_2 s_1 s_2;$$

$$(b3) \ s_2 s_6 s_2 = s_6 s_2 s_6;$$

$$(b4) \ s_6 s_1 s_2 s_6 = s_2 s_6 s_1 s_2.$$

(2) Under the presentation of G_{25} in (1), we have

$$(b5) \ s_1 s_6 s_1 = s_6 s_1 s_6.$$

Proof. (a2) is the same as (b2). By transition (5.4.1), relation (a1) (resp. (a3)) is equivalent to (b1) (resp. (b3)). Then under the assumption of relation (a3) (hence (b3)), relation (a4) is equivalent to (b4). So (1) follows by Proposition 5.3. Now we show (2). Left-multiplying s_6^{-1} and right-multiplying s_1 on both sides of (b4) and then applying (b2)-(b3), we get

$$(5.4.2) \quad s_1 s_2 s_6 s_1 = s_6^{-1} s_2 s_6 s_1 s_2 s_1 = s_2 s_6 s_2^{-1} s_1 s_2 s_1 = s_2 s_6 s_1 s_2.$$

This, together with (b4), implies that

$$(5.4.3) \quad s_1 s_2 s_6 s_1 = s_6 s_1 s_2 s_6.$$

Left-multiplying $s_1^{-1} s_2^{-1}$ on both sides of (5.4.2) and then applying (b2), we get

$$(5.4.4) \quad s_2 s_1^{-1} s_6 s_1 = s_1^{-1} s_6 s_1 s_2.$$

Then left-multiplying s_1^{-1} and right-multiplying s_6^{-1} on both sides of (5.4.3), we get

$$(5.4.5) \quad s_2 s_6 s_1 s_6^{-1} = s_1^{-1} s_6 s_1 s_2.$$

Therefore (b5) follows by (5.4.4) and (5.4.5). \square

With (5.4.3) instead of (b3) in the proposition, we get another congruent presentation of G_{25} .

Corollary. *The group G_{25} can be presented by the generator set $S_2 = \{s_1, s_2, s_6\}$ and the relations:*

$$\begin{aligned} (b1) \quad & s_1^3 = s_2^3 = s_6^3 = 1; \\ (b2) \quad & s_1 s_2 s_1 = s_2 s_1 s_2; \\ (b3') \quad & s_1 s_2 s_6 s_1 = s_2 s_6 s_1 s_2; \\ (b4) \quad & s_6 s_1 s_2 s_6 = s_2 s_6 s_1 s_2. \end{aligned}$$

The proof is left to the readers. Clearly, the presentation in the corollary is congruent to that in the proposition. With (b2) replaced by either (b3) or (b5) in the corollary, we get two more congruent presentations of G_{25} . One might notice that relations (b3'), (b4) are similar to (a2), (a3) in Proposition 3.3 for the group G_{12} with s_3 instead of s_6 . But it should be indicated that the orders of the generators for G_{25} are different from those for G_{12} .

§6. Simple root systems and presentations for the group G_{26} .

In this section, we do the same for the complex reflection group G_{26} . Unlike the previous three groups, the generators of G_{26} have two different orders, and the root system of G_{26} consists of two G_{26} -orbits. There are two inequivalent simple root systems and two non-congruent presentations for G_{26} (see 6.2 and Propositions 6.3, 6.4).

6.1. Let V , ϵ_i be as in 4.1. The vectors $e_1 = 2^{-1/2}(\epsilon_2 - \epsilon_3)$, $e_2 = \epsilon_3$ and $e_3 = 3^{-1/2}i(\epsilon_1 + \epsilon_2 + \epsilon_3)$ form a basis in V . Denote by (a, b, c) a vector $ae_1 + be_2 + ce_3$. Let $R_{26} = \{(-\omega)^j e_i \mid 1 \leq i \leq 21, j \in J\}$ be with

$$\begin{aligned} e_1 &= (1, 0, 0), & e_2 &= (0, 1, 0), & e_3 &= (0, 0, 1), & e_4 &= (1, y, 0), \\ e_5 &= (\sqrt{2}, 1, 0), & e_6 &= (0, 1, -\bar{\omega}), & e_7 &= (0, \bar{\omega}, 1), & e_8 &= (-\omega, y, 0), \\ e_9 &= (1, y, -\bar{\omega}y), & e_{10} &= (\sqrt{2}, 1, -\bar{\omega}), & e_{11} &= (\sqrt{2}, 1, \omega), & e_{12} &= (-\omega, y, -\bar{\omega}y), \\ e_{13} &= (1, y, \omega y), & e_{14} &= (\sqrt{2}, 1 - \omega, -\bar{\omega}), & e_{15} &= (\sqrt{2}, 2, \omega), & e_{16} &= (-\omega, y, \omega y), \\ e_{17} &= (1, 3/\sqrt{2}, \omega y), & e_{18} &= (-\sqrt{2}\omega, 1 - \omega, -\bar{\omega}), & e_{19} &= (\sqrt{2}, 2, -\bar{\omega}), & e_{20} &= (2, 3/\sqrt{2}, \omega y), \\ e_{21} &= (\sqrt{2}, 2, \sqrt{3}i), \end{aligned}$$

where $J = \{0, 1, 2, 3, 4, 5\}$ and $y = \frac{1}{\sqrt{2}}(1 - \omega)$. Let s_i be the reflection in V with respect to the vector e_i . Then the action of s_i on e_j , $i, j = 1, 2, 3$, are as follows.

e_i	$s_1(e_i)$	$s_2(e_i)$	$s_3(e_i)$
e_1	$-e_1$	e_4	e_1
e_2	e_5	ωe_2	e_6
e_3	e_3	e_7	ωe_3

The group generated by s_1, s_2, s_3 is G_{26} . R_{26} is the root system for G_{26} with $B_1 = \{e_1, e_2, e_3\}$ a simple root system (see [7]).

6.2. Now we consider all the inequivalent simple root systems B_k , $k \geq 1$. By Proposition 1.8, we have $|B_k| = 3$. The set R_{26} consists of two G_{26} -orbits: $R_1 = \{\omega^j e_i \mid i \in I, j \in J\}$ and $R_2 = \{\omega^j e_k \mid j \in J, k \in K\}$, where $I = \{1, 4, 8, 9, 12, 13, 16, 17, 20\}$ and $K = \{2, 3, 5, 6, 7, 10, 11, 14, 15, 18, 19, 21\}$. By Lemma 1.3 and condition $G_{26}B_k = R_{26}$, we have $B_k \cap R_i \neq \emptyset$ for $i = 1, 2$. We may assume $e_1 \in B_k$ up to G_{26} -action. By the action of the s_i 's on R_{26} , we see that each B_k contains one root in R_1 and two roots in R_2 . All the inequivalent simple root systems for R_{26} are as follows.

$$B_1 = \{e_1, e_2, e_3\}, B_2 = \{e_1, e_2, e_6\}.$$

We have $\mathbf{n}(B_1) = (G(3, 1, 2), G_4, \mathbb{Z}_2 \times \mathbb{Z}_3)$ and $\mathbf{n}(B_2) = (G(3, 1, 2), G(3, 1, 2), G_4)$ (see 1.10 for $\mathbf{n}(B)$ and [5] for $G(3, 1, 2)$).

6.3. Next we consider all the non-congruent presentations of G_{26} by generators and relations according to the result in 6.2.

Proposition. *The group G_{26} can be presented by the generator set $S_1 = \{s_1, s_2, s_3\}$ and the relations:*

$$(a1) \ s_1^2 = s_2^3 = s_3^3 = 1;$$

$$(a2) \ s_1 s_2 s_1 s_2 = s_2 s_1 s_2 s_1;$$

$$(a3) \ s_2 s_3 s_2 = s_3 s_2 s_3;$$

$$(a4) \ s_1 s_3 = s_3 s_1.$$

Proof. This follows by [1, Table 1]. \square

6.4. We have

$$(6.4.1) \quad s_6 = s_2^{-1} s_3 s_2 \quad \text{and} \quad s_3 = s_2 s_6 s_2^{-1}.$$

Proposition. *(1) The group G_{26} can be presented by the generator set $S_2 = \{s_1, s_2, s_6\}$ and the relations:*

$$(b1) \ s_1^2 = s_2^3 = s_6^3 = 1;$$

$$(b2) \ s_1 s_2 s_1 s_2 = s_2 s_1 s_2 s_1;$$

$$(b3) \ s_2 s_6 s_2 = s_6 s_2 s_6;$$

$$(b4) \ s_2 s_6 s_1 s_2 = s_6 s_1 s_2 s_6.$$

(2) Under the presentation of G_{26} in (1), we have

$$(b5) \ s_1 s_6 s_1 s_6 = s_6 s_1 s_6 s_1.$$

Proof. (1) Relations (a2) and (b2) are the same. By transition (6.4.1), relation (a1) (resp., (a3)) is equivalent to (b1) (resp., (b3)). Then under the assumption of relations (6.4.1) and (b3) (hence (a3)), we have

$$(a4) \iff s_1 \cdot s_2 s_6 s_2^{-1} = s_2 s_6 s_2^{-1} \cdot s_1 \iff s_1 \cdot s_2 s_6 s_2^{-1} = s_6^{-1} s_2 s_6 \cdot s_1 \iff (b4)$$

This implies (1). Now we show (2). Rewrite (b4) as

$$(6.4.2) \quad s_2^{-1} \cdot s_6 s_1 \cdot s_2 = s_6 \cdot s_1 s_2 \cdot s_6^{-1}.$$

Squaring both sides and then applying (b2), we get

$$(6.4.3) \quad s_2^{-1} \cdot s_6 s_1 s_6 s_1 \cdot s_2 = s_6 \cdot s_2 s_1 s_2 s_1 \cdot s_6^{-1}.$$

Left-multiplying $s_2^{-1} s_6^{-1} s_2$ and right-multiplying $s_6 s_2 s_6^{-1}$ on both sides of (6.4.3) and then applying (b3), we get

$$(6.4.4) \quad s_2^{-1} \cdot s_1 s_6 s_1 s_6 \cdot s_2 = s_6 \cdot s_1 s_2 s_1 s_2 \cdot s_6^{-1}$$

So (b5) follows by (b2), (6.4.3) and (6.4.4). \square

§7. Essential presentations.

7.1. According to the definition in 1.9, if (S, \mathcal{P}) is a presentation of a reflection group G , and if \mathcal{P}' is a set of relations on S , including \mathcal{P} as a subset, then (S, \mathcal{P}') is also a presentation of G .

A presentation (S, \mathcal{P}) of G is *essential* if (S, \mathcal{P}_0) is not a presentation of G for any proper subset \mathcal{P}_0 of \mathcal{P} .

Checking the essentiality for a presentation of a group is usually a subtle task.

In a presentation (S, \mathcal{P}) of G , a relation $P \in \mathcal{P}$ is *redundant* if P is a consequence of the relations in $\mathcal{P} \setminus \{P\}$. Thus a presentation (S, \mathcal{P}) of G is essential if and only if \mathcal{P} contains no redundant relation.

Let (S, \mathcal{P}) be one of the presentations we get so far for the groups $G = G_{12}, G_{24}, G_{25}, G_{26}$. We define \mathcal{P}^{re} , which is obtained from \mathcal{P} in the following way: Let B be a simple root system for G such that $S = \{s_\alpha \mid \alpha \in B\}$. For any $\alpha \in B$, if $G\alpha \cap B$ contains $k > 1$ elements, then we remove any $k - 1$ of the relations $s_\beta^{o(s_\beta)} = 1, \beta \in G\alpha \cap B$, from \mathcal{P} .

Example 7.2. (1) Let (S, \mathcal{P}) be the presentation of G_{12} given in Proposition 3.3. Then \mathcal{P}^{re} is obtained from \mathcal{P} by removing relations $s_2^2 = 1$ and $s_3^2 = 1$.

(2) Let (S, \mathcal{P}) be the presentation of G_{26} in Proposition 6.3. Then \mathcal{P}^{re} is obtained from \mathcal{P} by removing relation $s_3^3 = 1$.

The main result of the present section is as follows.

Theorem 7.3. *For any presentation (S, \mathcal{P}) we get so far for the group $G = G_{12}, G_{24}, G_{25}$ or G_{26} , the above defined $(S, \mathcal{P}^{\text{re}})$ is again a presentation of G . Moreover, $(S, \mathcal{P}^{\text{re}})$ is essential.*

We shall show the theorem in the remaining part of the section. The following is a simple fact in the group theory.

Lemma 7.4. *In a group G , if $x, y, z \in G$ satisfy $xy = yz$, then $o(x) = o(z)$.*

Then the conclusion that $(S, \mathcal{P}^{\text{re}})$ is a presentation of the group G follows by Lemma 7.4. In Example 7.2 (1), relations $s_2^2 = s_3^2 = 1$ are the consequence of relations $s_1^2 = 1, s_1 \cdot s_2 s_3 s_1 = s_2 s_3 s_1 \cdot s_2$ and $s_3 \cdot s_1 s_2 s_3 = s_1 s_2 s_3 \cdot s_1$ in \mathcal{P}^{re} (see Proposition 3.3, (a1)-(a3)). In Example 7.2 (2), relation $s_3^3 = 1$ is a consequence of relations $s_2^3 = 1$ and $s_2 \cdot s_3 s_2 = s_3 s_2 \cdot s_3$ in \mathcal{P}^{re} (see Proposition 6.3, (a1)-(a4)). So in each of these two examples, $(S, \mathcal{P}^{\text{re}})$ is a presentation of the respective group. All the other cases could be checked similarly. Thus it makes sense to call $(S, \mathcal{P}^{\text{re}})$ the *reduced form* for the presentation (S, \mathcal{P}) .

Let $(x1^{\text{re}})$ be the set of relation(s) in the presentation $(S, \mathcal{P}^{\text{re}})$ obtained from $(x1)$ in the presentation (S, \mathcal{P}) by removing certain relation(s) for $x=a, b, c, \dots$. For example, when (S, \mathcal{P}) is the presentation of the group G_{12} (resp., G_{26}) in Proposition 3.3 (resp., Proposition 6.3), $(a1^{\text{re}})$ in $(S, \mathcal{P}^{\text{re}})$ contains a single relation $s_1^2 = 1$ (resp., two relations $s_1^2 = 1$ and $s_2^3 = 1$).

The following fact is useful in checking the essentiality for a presentation of a complex reflection group.

Lemma 7.5. *Let G be a complex reflection group with (S, \mathcal{P}) its presentation.*

(1) *The relation set \mathcal{P} contains at least one relation which either involves more than two generators, or has the form $s^m = 1$ with $m > 2$ for some $s \in S$.*

(2) The relation(s) contained in $(x1^{\text{re}})$, $x=a,b,\dots$, is(are) not redundant in the reduced form of (S, \mathcal{P}) when (S, \mathcal{P}) is one of those we get in the previous sections.

(3) Let (S, \mathcal{P}') be another presentation of G with $\mathcal{P}' \subseteq \mathcal{P}$. For any $P \in \mathcal{P}'$, if P is redundant in \mathcal{P}' , then P is redundant in \mathcal{P} .

Proof. (1) is true since a complex reflection group is not a Coxeter group. Then (2) follows by the following fact: substituting $s_i = c$ for all $1 \leq i \leq 3$ and any fixed $c \in \mathbb{C} \setminus \{0\}$, all the relations in $\mathcal{P} \setminus \{P \mid P \in (x1)\}$ remain valid. Finally, as a consequence of relations in $\mathcal{P}' \setminus \{P\}$, P is clearly a consequence of relations in $\mathcal{P} \setminus \{P\}$. So we get (3). \square

Thus it remains to show the essentiality of the reduced form $(S, \mathcal{P}^{\text{re}})$ for any presentation (S, \mathcal{P}) we get so far for the group $G = G_{12}, G_{24}, G_{25}$ or G_{26} . The following result will be used for this purpose.

Lemma 7.6. *Suppose that a reflection group G has two presentations (S, \mathcal{P}) and (S', \mathcal{P}') , where $\mathcal{P} = \{P_i \mid i \in I\}$ and $\mathcal{P}' = \{P'_i \mid i \in I'\}$ are relation sets with (I, \leq_I) , $(I', \leq_{I'})$ finite posets (the word “poset” means “partially ordered set”). Suppose that there exists an order-preserving surjection $\phi : (I, \leq_I) \rightarrow (I', \leq_{I'})$ such that for any $k \in I$, relation set $\{P_i \mid i \leq_I k\}$ is equivalent to $\{P'_{i'} \mid i' \leq_{I'} \phi(k)\}$. Suppose that $m, \phi(m)$ are maximal elements in I, I' , respectively with $\phi^{-1}(\phi(m)) = \{m\}$. Then relation P_m is not redundant in \mathcal{P} if and only if $P'_{\phi(m)}$ is not redundant in \mathcal{P}' .*

Proof. First show the implication “ \implies ”. We argue by contrary. Suppose that $P'_{\phi(m)}$ is a consequence of all the other relations in \mathcal{P}' . By our assumption, we see that relation P_m is a consequence of the relations in $\{P'_i \mid i \leq_{I'} \phi(m)\}$ and hence a consequence of the relations in $\mathcal{P}' \setminus \{P'_{\phi(m)}\}$. Since relation set $\mathcal{P} \setminus \{P_m\}$ is equivalent to relation set $\mathcal{P}' \setminus \{P'_{\phi(m)}\}$ by our condition, this implies that P_m is a consequence of the relations in $\mathcal{P} \setminus \{P_m\}$, a contradiction.

The implication “ \impliedby ” can be shown similarly by interchanging the roles of m, I, \mathcal{P} with $\phi(m), I', \mathcal{P}'$ respectively. \square

7.7. By Lemma 7.5 (2), we see that relation $(x1^{\text{re}})$, $x=a,b,\dots$, is not redundant in the respective reduced presentation for the group $G = G_{12}, G_{24}, G_{25}$ or G_{26} . In the following discussion, when we talk about the redundancy of a relation P in a (or reduced) presentation (S, \mathcal{P}) (or $(S, \mathcal{P}^{\text{re}})$) of G , we often simply state “ P is not redundant” for “ P is not redundant in (S, \mathcal{P}) (or $(S, \mathcal{P}^{\text{re}})$)” if (S, \mathcal{P}) (or $(S, \mathcal{P}^{\text{re}})$) is clear in the context. By Lemma 7.5 (3),

when a relation P is in \mathcal{P}^{re} , to show that P is not redundant in \mathcal{P}^{re} , we need only show that P is not redundant in \mathcal{P} (we always do in this way in the subsequent discussion).

7.8. Let us first consider the group G_{24} . We can show that the generators s_1, s_2 are symmetric for the presentation of G_{24} in Proposition 4.3 in the following sense: suppose that relations (ai') , $1 \leq i \leq 3$, are obtained from (ai) by interchanging s_1, s_2 , then two relation sets $(a1)-(a3)$ and $(a1')-(a3')$ can be obtained from one to another. We can also show that s_1, s_3 (resp. s_1, s_3, s_8) are symmetric for the presentation of G_{24} in Proposition 4.4 (resp. Proposition 4.5). Since the reflection group G_{24} is complex, relations $(a5)$, $(b5)$ are not redundant by Lemma 7.5 (1), and neither are relations $(a4)$ and $(b4)$ by the proof of Proposition 4.4 and by Lemma 7.6. By symmetry, relations $(a2)$ and $(b3)$ are not redundant. Relations $(a2)$, $(b2)$ and $(c2)$ are the same. So neither of $(b2)$ and $(c2)$ is redundant by the proof of Propositions 4.4, 4.5 and by Lemma 7.6. Again by the proof of Proposition 4.5 and by Lemma 7.6, relation $(c5)$ is not redundant. By the symmetry of s_1, s_8 for the presentation of G_{24} in Proposition 4.5, neither of relations $(c3)$ and $(c4)$ is redundant. Finally, by the proof of Proposition 4.5 and by Lemma 7.6, relation $(a3)$ is not redundant. This proves the essentiality of the reduced forms for all the presentations of G_{24} in Propositions 4.3, 4.4 and 4.5.

7.9. Now we consider the group G_{25} . We claim that relation $(a2)$ is not redundant. For, relation $s_1 s_2 = s_2 s_1$ together with relations $(a1)$, $(a3)$ and $(a4)$ determine a reducible reflection group $\langle s_1 \rangle \times \langle s_2, s_3 \rangle$, meaning that $(a2)$ is not a consequence of relations $(a1)$, $(a3)$, $(a4)$. Similarly, $(a3)$ is not redundant. We claim that relation $(a4)$ is not redundant. For, by replacing s_6 by s_3 for the presentation of G_{25} in Proposition 5.4, relations $(b1)-(b3)$ becomes $(a1)-(a3)$, respectively, and $(b5)$ becomes $s_1 s_3 s_1 = s_3 s_1 s_3$, differing from $(a4)$, which shows that relation $(a4)$ is not a consequence of relations $(a1)-(a3)$. So the reduced form for the presentation of G_{25} in Proposition 5.3 is essential. Then by the proof of Proposition 5.4 and by Lemma 7.6, we see that neither of relations $(b2)$ and $(b4)$ is redundant. Then by the symmetry of s_1, s_2, s_6 for the presentation of G_{25} in Proposition 5.4, we see that relation $(b3)$ is not redundant, too. This implies that the reduced form for the presentation of G_{25} in Proposition 5.4 is essential. Finally, to show that the reduced form for the presentation of G_{25} in Corollary 5.5 is essential, we need only show that relation $(b3')$ is not redundant. But this follows by the symmetry of

s_1, s_2, s_6 for the presentation of G_{25} in Corollary 5.5.

The proof is similar for the essentiality of the reduced form for the presentations of the group G_{26} in Propositions 6.3 and 6.4.

7.10. Finally we consider the group G_{12} . Relation (a3) is not redundant, since (b1) and (b4) become (a1) and (a2) respectively by replacing s_{10} by s_3 , while (b3) does not become (a6) under the same substitution. By symmetry of s_1, s_2, s_3 for the presentation of G_{12} in Proposition 3.3, we see that relation (a2) is not redundant, too. So the reduced form for the presentation of G_{12} in Proposition 3.3 is essential. By Lemma 7.5 (1), relation (b4) is not redundant. We claim that relation (b3) is not redundant. For, relation $s_1 s_{10} s_1 s_{10} s_1 s_{10} = s_{10} s_1 s_{10} s_1 s_{10} s_1$ does not conflict with (b1), (b2) and (b4) by observing the presentation of G_{12} in Proposition 3.3 with s_3 in the place of s_{10} . Next we claim that relation (b2) is not redundant. To see this, one need only notice that with respect to the presentation of G_{12} in Proposition 3.3, the images $\bar{s}_1, \bar{s}_2, \bar{s}_3$ of the generators s_1, s_2, s_3 in the quotient group of G_{12} over its center Z (Z has order two) satisfy relations (b1), (b3), (b4) with $\bar{s}_1, \bar{s}_2, \bar{s}_3$ in the places of s_1, s_2, s_{10} respectively, and relation $\bar{s}_1 \bar{s}_2 \bar{s}_1 = \bar{s}_2 \bar{s}_1 \bar{s}_2$. Hence the reduced form for the presentation of G_{12} in Proposition 3.4 is essential. The non-redundancy of relation (c3) follows by the proof of Proposition 3.5 and by Lemma 7.6. Then by the symmetry of s_1, s_7 for the presentation of G_{12} in Proposition 3.5, we see that relation (c2) is not redundant, too. This implies that the reduced form for the presentation of G_{12} in Proposition 3.5 is essential. By the proof of Proposition 3.6 and by Lemma 7.6, we see that neither of relations (d2) and (d3) is redundant and hence the reduced form for the presentation of G_{12} in Proposition 3.6 is essential. Finally consider the presentation of G_{12} in Proposition 3.7. Relation (e5) is not redundant by Lemma 7.5 (1). Relation (e2) is not redundant since one gets a presentation of the quotient group G_{12}/Z (isomorphic to the symmetric group S_4) by replacing (e2) by $s_1 s_7 = s_7 s_1$ in Proposition 3.7. Next we claim that relation (e3) is not redundant. For, substituting s_1, s_7, s_8 by s_2, s_8, s_1 in relations (e1)-(e5) respectively, followed by replacing a resulting relation $s_1 s_2 s_1 = s_2 s_1 s_2$ (which comes from (e3)) by (d6), all the relations we get are the consequence of the presentation of G_{12} in Proposition 3.6. This shows the claim. By symmetry of s_1, s_7 in the presentation, we see that relation (e4) is not redundant, too. So the reduced form for the presentation of G_{12} in Proposition 3.7 is essential.

Appendix.

Here we record the results of L. Wang and P. Zeng on the numbers $N(G)$ of inequivalent

simple root systems for the primitive complex reflection groups $G_7, G_{11}, G_{13}, G_{15}, G_{19}, G_{22}, G_{27}, G_{29}, G_{31}, G_{32}, G_{33}$.

G	$N(G)$	G	$N(G)$
G_7	2	G_{11}	4
G_{13}	4	G_{15}	4
G_{19}	6	G_{22}	18
G_{27}	6	G_{29}	9
G_{31}	61	G_{32}	5
G_{33}	14		

Since any complex reflection group generated by one or two reflections has a unique inequivalent simple root system, so far G_{34} is the only group left unknown concerning the inequivalent simple root systems.

REFERENCES

1. M. Broue, G. Malle and R. Rouquier, *Complex reflection groups, Braid groups, Hecke algebras*, J. Reine Angew. Math. **500** (1998), 127–190.
2. Arjeh M. Cohen, *Finite complex reflection groups*, Ann. scient. Éc. Norm. Sup. 4^e série t. **9** (1976), 379-436.
3. R. B. Howlett and J. Y. Shi, *On regularity of finite reflection groups*, manuscripta mathematica **102(3)** (2000), 325-333.
4. D. W. Koster, *Complex reflection groups*, Ph.D. thesis, University of Wisconsin (1975).
5. G. C. Shephard and J. A. Todd, *Finite unitary reflection groups*, Canad. J. Math. **6** (1954), 274-304.
6. J. Y. Shi, *Certain imprimitive reflection groups and their generic versions*, Trans. Amer. Math. Soc. **354** (2002), 2115-2129.
7. V. L. Popov, *Discrete complex reflection groups*, Communications of the Mathematical Institute, Rijksuniversiteit Utrecht **15**. (1982), 1-89.
8. Li Wang, *Simple root systems and some presentations for the primitive reflection groups generated by reflections of order 2*, Master Dissertation, East China Normal University, 2003.
9. Peng Zeng, *Simple root systems and presentations for the primitive reflection groups containing a reflection of order > 2* , Master Dissertation, East China Normal University, 2003.

Email: jyshi@math.ecnu.edu.cn