

## Pronunciation of mathematical expressions

The pronunciations of the most common mathematical expressions are given in the list below. In general, the shortest versions are preferred (unless greater precision is necessary).

### 1. Logic

$\exists$	there exists
$\forall$	for all
$p \Rightarrow q$	$p$ implies $q$ / if $p$ , then $q$
$p \Leftrightarrow q$	$p$ if and only if $q$ / $p$ is equivalent to $q$ / $p$ and $q$ are equivalent

### 2. Sets

$x \in A$	$x$ belongs to $A$ / $x$ is an element (or a member) of $A$
$x \notin A$	$x$ does not belong to $A$ / $x$ is not an element (or a member) of $A$
$A \subset B$	$A$ is contained in $B$ / $A$ is a subset of $B$
$A \supset B$	$A$ contains $B$ / $B$ is a subset of $A$
$A \cap B$	$A$ cap $B$ / $A$ meet $B$ / $A$ intersection $B$
$A \cup B$	$A$ cup $B$ / $A$ join $B$ / $A$ union $B$
$A \setminus B$	$A$ minus $B$ / the difference between $A$ and $B$
$A \times B$	$A$ cross $B$ / the cartesian product of $A$ and $B$

### 3. Real numbers

$x + 1$	$x$ plus one
$x - 1$	$x$ minus one
$x \pm 1$	$x$ plus or minus one
$xy$	$xy$ / $x$ multiplied by $y$
$(x - y)(x + y)$	$x$ minus $y$ , $x$ plus $y$
$\frac{x}{y}$	$x$ over $y$
$=$	the equals sign
$x = 5$	$x$ equals 5 / $x$ is equal to 5
$x \neq 5$	$x$ (is) not equal to 5

$x \equiv y$	$x$ is equivalent to (or identical with) $y$
$x \not\equiv y$	$x$ is not equivalent to (or identical with) $y$
$x > y$	$x$ is greater than $y$
$x \geq y$	$x$ is greater than or equal to $y$
$x < y$	$x$ is less than $y$
$x \leq y$	$x$ is less than or equal to $y$
$0 < x < 1$	zero is less than $x$ is less than 1
$0 \leq x \leq 1$	zero is less than or equal to $x$ is less than or equal to 1
$ x $	mod $x$ / modulus $x$
$x^2$	$x$ squared / $x$ (raised) to the power 2
$x^3$	$x$ cubed
$x^4$	$x$ to the fourth / $x$ to the power four
$x^n$	$x$ to the $n$ th / $x$ to the power $n$
$x^{-n}$	$x$ to the (power) minus $n$
$\sqrt{x}$	(square) root $x$ / the square root of $x$
$\sqrt[3]{x}$	cube root (of) $x$
$\sqrt[4]{x}$	fourth root (of) $x$
$\sqrt[n]{x}$	$n$ th root (of) $x$
$(x + y)^2$	$x$ plus $y$ all squared
$\left(\frac{x}{y}\right)^2$	$x$ over $y$ all squared
$n!$	$n$ factorial
$\hat{x}$	$x$ hat
$\bar{x}$	$x$ bar
$\tilde{x}$	$x$ tilde
$x_i$	$x_i$ / $x$ subscript $i$ / $x$ suffix $i$ / $x$ sub $i$
$\sum_{i=1}^n a_i$	the sum from $i$ equals one to $n$ $a_i$ / the sum as $i$ runs from 1 to $n$ of the $a_i$

#### 4. Linear algebra

$\ x\ $	the norm (or modulus) of $x$
$\overrightarrow{OA}$	$OA$ / vector $OA$
$\overline{OA}$	$OA$ / the length of the segment $OA$
$A^T$	$A$ transpose / the transpose of $A$
$A^{-1}$	$A$ inverse / the inverse of $A$

## 5. Functions

$f(x)$	$fx$ / $f$ of $x$ / the function $f$ of $x$
$f : S \rightarrow T$	a function $f$ from $S$ to $T$
$x \mapsto y$	$x$ maps to $y$ / $x$ is sent (or mapped) to $y$
$f'(x)$	$f$ prime $x$ / $f$ dash $x$ / the (first) derivative of $f$ with respect to $x$
$f''(x)$	$f$ double-prime $x$ / $f$ double-dash $x$ / the second derivative of $f$ with respect to $x$
$f'''(x)$	$f$ triple-prime $x$ / $f$ triple-dash $x$ / the third derivative of $f$ with respect to $x$
$f^{(4)}(x)$	$f$ four $x$ / the fourth derivative of $f$ with respect to $x$
$\frac{\partial f}{\partial x_1}$	the partial (derivative) of $f$ with respect to $x_1$
$\frac{\partial^2 f}{\partial x_1^2}$	the second partial (derivative) of $f$ with respect to $x_1$
$\int_0^\infty$	the integral from zero to infinity
$\lim_{x \rightarrow 0}$	the limit as $x$ approaches zero
$\lim_{x \rightarrow +0}$	the limit as $x$ approaches zero from above
$\lim_{x \rightarrow -0}$	the limit as $x$ approaches zero from below
$\log_e y$	$\log y$ to the base $e$ / $\log$ to the base $e$ of $y$ / natural $\log$ (of) $y$
$\ln y$	$\log y$ to the base $e$ / $\log$ to the base $e$ of $y$ / natural $\log$ (of) $y$

Individual mathematicians often have their own way of pronouncing mathematical expressions and in many cases there is no generally accepted “correct” pronunciation.

Distinctions made in writing are often not made explicit in speech; thus the sounds  $fx$  may be interpreted as any of:  $fx$ ,  $f(x)$ ,  $f_x$ ,  $FX$ ,  $\overline{FX}$ ,  $\overrightarrow{FX}$ . The difference is usually made clear by the context; it is only when confusion may occur, or where he/she wishes to emphasise the point, that the mathematician will use the longer forms:  $f$  multiplied by  $x$ , the function  $f$  of  $x$ ,  $f$  subscript  $x$ , line  $FX$ , the length of the segment  $FX$ , vector  $FX$ .

Similarly, a mathematician is unlikely to make any distinction in speech (except sometimes a difference in intonation or length of pauses) between pairs such as the following:

$$\begin{aligned}
 x + (y + z) & \quad \text{and} \quad (x + y) + z \\
 \sqrt{ax} + b & \quad \text{and} \quad \sqrt{ax + b} \\
 a^n - 1 & \quad \text{and} \quad a^{n-1}
 \end{aligned}$$

The primary reference has been David Hall with Tim Bowyer, *Nucleus, English for Science and Technology, Mathematics*, Longman 1980. Glen Anderson and Matti Vuorinen have given good comments and supplements.

# Reading Mathematical Expressions

(A. Oancea, S. Maillot, C. Mitschi)

**Note:** Some groups of letters are underlined in order to draw one's attention to their pronunciation.

## Basics

$a + b$	$a$ plus $b$
$a - b$	$a$ <u>minus</u> $b$
$a \cdot b$	$ab$ , $a$ times $b$
$\frac{a}{b}$ , $a/b$	$a$ over $b$ , $a$ divided by $b$
$\frac{1}{2}$ , $\frac{1}{3}$ , $\frac{1}{4}$ , $\dots$ , $\frac{1}{10}$	one half, one third, one fourth, ... , one tenth
$\frac{5}{2}$ , $\frac{2}{3}$ , $\dots$ , $\frac{7}{10}$	five halves, two thirds, ... , seven tenths
$a = b$	$a$ equals <u><math>b</math></u> , $a$ is equal to $b$
$a \neq b$	$a$ different from $b$ , $a$ not equal to $b$
$a < b$	$a$ (strictly) less than $b$
$a \leq b$	$a$ less than or equal to $b$
$a > b$	$a$ (strictly) bigger than $b$ , $a$ greater than $b$
$a \geq b$	$a$ greater than or equal to $b$

## Powers and roots

$a^b$	$a$ to the $b$ ,
	$a$ to the $b$ -th (power) [if $b$ is a positive integer]
$x^2$	$x$ squared
$x^3$	$x$ cubed
$x^{-1}$	$x$ inverse
$\sqrt[n]{t}$	$n$ -th root of $t$
$\sqrt{t}$	square root of $t$
$\sqrt[3]{t}$	cubic root of $t$

## Sets

$\emptyset$	(the) empty set
$A \cup B$	$A$ union $B$
$A \cap B$	$A$ intersected with $B$
$A^c$	the complement of $A$
$A \setminus B$	$A$ minus $B$
$A \times B$	$A$ times $B$
$x \in A$	$x$ in $A$ , $x$ belongs to $A$ , $x$ belonging to $A$

## Miscellaneous

5%	five percent
30°	thirty degrees
$x_k$	$x$ $k$
$x_i^j$	$x$ $i$ $j$ [if $j$ is an index, not an exponent!]
$\sum_{k=1}^n k^2$	sum $k$ equals 1 to $n$ of $k^2$ , sum for $k$ (running) from 1 to $n$ of $k^2$ , summation $k$ from 1 to $n$ of $k^2$
$\prod_{k=1}^n \frac{2k+1}{2k+2}$	product $k$ equals 1 to $n$ of $2k + 1$ over $2k + 2$ product for $k$ (running) from 1 to $n$ of $2k + 1$ over $2k + 2$
$n!$	$n$ factorial
partie entière de $x$	integer part of $x$
$ x $	absolute value of $x$ (if $x$ is a real number)
$ z $	modulus of $z$ (if $z$ is a complex number)
$\operatorname{Re}(z)$ , $\operatorname{Im}(z)$	real part of $z$ , imaginary part of $z$
$\ x\ $	norm of $x$
$\langle v, w \rangle$	scalar product of $v$ and $w$
cos sin tan etc.	cos <u>i</u> ne/cosinus s <u>i</u> ne/sinus tangent etc.
$\eta$ $\theta$ $\xi$	<u>e</u> ta [íta] <u>t</u> heta [thíta] <u>x</u> i [ksái]
$\pi$ $\sigma$ $\chi$ $\psi$	<u>p</u> i [pái] <u>s</u> igma [zígamma] <u>ch</u> i [kái] <u>ps</u> i [sái]
$\mathbb{R}^2$ , $\mathbb{R}^3$ , $\mathbb{R}^n$	$\mathbb{R}$ 2, $\mathbb{R}$ 3, $\mathbb{R}$ $n$
$(blablabla) \cdot (blbl)$	<i>blablabla</i> , the whole times <i>blbl</i>
$\frac{blablabla}{blbl}$	<i>blablabla</i> , the whole divided by <i>blbl</i>
$x_1, \dots, x_n$	$x_1$ up to $x_n$

## Calculus

$f'$	$f$ prime, $f$ dashed
$\frac{d}{dx}$	$d$ by $dx$
$\frac{df}{dx}, \frac{\partial f}{\partial x}$	$df$ by $dx$
$\partial_x f$	$d_x f$ , partial derivative of $f$ with respect to $x$
$\int_a^b f(s) ds$	integral from $a$ to $b$ (of) $f(s) ds$
$\iint_D, \iiint_D$	double integral, triple integral over the domain $D$
$\pm\infty$	plus/minus infinity
$\lim_{x \rightarrow a} f(x)$	(the) limit of $f(x)$ as $x$ tends/goes to $a$ , (the) limit of $f$ of $x$ as $x$ tends/goes to $a$
$\log(x), \log_a x$	logarith <u>m</u> of $x$ , logarith <u>m</u> in base $a$ of $x$
$\exp(x), e^x$	exponential of $x$ , $e$ to the $x$

## Functions

$f : U \rightarrow V$	$f$ from $U$ to $V$
$f(x)$	$f$ of $x$
$x \mapsto f(x)$	$x$ maps to $f(x)$
of class $C^k$	of class $C^k$
of class $C^\infty$	of class $C$ infinity
the Lebesgue spaces $L^p, L^\infty$	the Lebesgue spaces $L^p, L$ infinity
the Sobolev spaces $H^k, W^{k,p}$	the <u>S</u> obolev spaces $H^k, W^k p$